



Royal  
Horticultural  
Society

Sharing the best in Gardening

# The International Rhododendron Register & Checklist (2004)

*Eleventh Supplement*



© 2016 The Royal Horticultural Society  
80 Vincent Square, London SW1P 2PE, United Kingdom  
[www.rhs.org.uk](http://www.rhs.org.uk)

International Rhododendron Registrar:  
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057724

Printed and bound in the UK by Page Bros, Norwich

The previous supplement (Tenth Supplement) was published on 23 July 2015

Cover: *Rhododendron* 'Fireball', by Elsie Katherine Kohnlein  
Image courtesy of RHS Lindley Library

# The International Rhododendron Register and Checklist (2004)

## *Eleventh Supplement*

Names registered, January–December 2015	page 2
Corrections	page 17
Names and addresses	page 18

## Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first ten Supplements included names registered from 1 January 2003 to 31 December 2014. This eleventh Supplement includes all those registered in 2015.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction.

It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea
(az)	azaleodendron
cv.	cultivar
G	grown to first flower by
gp	Group
H	hybridised by
I	introduced by
N	named by
(r)	rhododendron
R	raised by
REG	registered by
S	selected by
(s)	seed parent
(v)	vireya

Where possible colour references are given to the *RHS Colour Chart* (1966 (reprinted 1986), 1995, 2001, 2007, 2015) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website ([www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform](http://www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform)) or from one of the national or regional registrars as follows:

**North America** Mr Michael Martin Mills, 632 Burnham Road, Philadelphia, Pennsylvania PA19119-3510, USA

**Australia** Mrs Lesley Eaton, PO Box 597, Snug, Tasmania 7054, Australia

**New Zealand** Mr Brian J. Coker, Stonecrop, 8B Barnsley Crescent, West Melton, New Zealand 7618

**Japan** Dr Katsuhisa Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

**Dr A.C. Leslie, International Rhododendron Registrar, 109 York Street, Cambridge CB1 2PZ, UK**  
(email: [alanleslie@rhs.org.uk](mailto:alanleslie@rhs.org.uk))

# Names registered January-December 2015

## (a) 'Adalmina'

cv. Deciduous azalea: 'Directeur Charles Baumann' (s) × Choptank River Hybrid × 'Maori'.

**H:** A. Väinölä (1989), **G:** University of Helsinki & Arboretum Mustila, **N:** A. Väinölä (2008), **I:** MTT Agrifood Research Finland (2010), **REG:** University of Helsinki (2015).

Fls 8–10/truss, narrowly funnel-shaped, 45–51 × 53–56mm, with 5 flat to wavy-edged lobes, buds strong red (53C–D), opening inside strong red (51A), deep pink (52B) and strong to deep purplish red (54A, 55A); outside deep pink (52B), deep purplish pink (54B) and strong pink (54C); dorsal lobe with a central vivid yellow (17B) blotch, with a thin red midvein; stamens and style vivid red (52A) to strong red (53C). Calyx 1–2mm long. Lvs broadly elliptic, 56–63 × 25–31mm, cuneate at base, acute to acuminate at apex, margins flat to decurved, matt green above, hairy. Shrub 1.5 × 1.8m in 20 years. Late June. Hardy.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119839).

Etymology: named for a princess in a Finnish fairytale written by Z. Topelius in 1893. First published in *Sorbifolia* 43(1):34–38 (2012).

## (r) 'Aldonis Vēriņš'

cv. Elepidote rhododendron: 'Nova Zembla' × unknown.

**H:** R. Kondratovičs (1987), **G:** R. Kondratovičs (1993), **N:** R. Kondratovičs (2015), **REG:** Rhododendron Nursery Babite, University of Latvia (2015).

Fls 22/strongly domed truss, funnel-shaped, 50 × 70mm, with 5 wavy-edged lobes, buds deep purplish red (61A), opening inside strong purplish red (71C), with greyish purple (N77A) spots on the dorsal lobes, outside vivid purplish red (67B); filaments red-purple, anthers white; style red-purple, with a black stigma. Calyx 1mm long, red-purple. Lvs elliptic, 130 × 70mm, rounded at base, pointed at apex, margins slightly wavy, matt green above, hairless. Bushy shrub, 1.7 × 1.7m in 10 years. Late May to early June. Hardy to –30C.

Nomenclatural standard: colour print provided by the registrant (WSY0119768). Etymology: named for a Latvian famous for his selections of iris, gladiolus and peony cultivars.

## (r) 'Atlantic Gold'

cv. Elepidote rhododendron: 'Percy Wiseman' (s) × ('Golden Star' × *wardii* (KW 4170)) × ((Jalisco Group × *yakushimanum*) × 'Odee Wright').

**H:** R.A. Furman (1994), **G:** R.A. Furman (2002), **N:** A. Furman (2015), **REG:** J. Delano (2015).

Fls 13/domed truss, funnel-shaped, 51 × 90mm, with 7 rounded, wavy-edged lobes, pale purplish pink (56B–C) in bud, becoming light yellow (10C) just before opening, opening inside brilliant greenish

yellow (6C) to light greenish yellow (6D), with deep red (53A) basal flares corresponding to each lobe (longer on the dorsal side), forming a seven point basal star 20mm in diameter, and with some green spotting beyond; outside light yellow (10C); stigma deep red (53A); not scented. Calyx 8mm long, light yellow (10B). Lvs elliptic, 140 × 56mm, oblique at base, obtuse at apex, margins flat, with a depressed central vein, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.7m in 20 years. Midseason (mid May on Cape Cod, Massachusetts, USA). Hardy to –21C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119841).

## (r) 'Babites Aiga'

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' × unknown.

**H:** R. Kondratovičs (1971), **G:** R. Kondratovičs (1977), **N:** R. Kondratovičs (2015), **I:** Rhododendron Nursery Babite, University of Latvia (2015), **REG:** Rhododendron Nursery Babite, University of Latvia (2015).

Fls 20/strongly domed truss, open funnel-shaped, 80 × 75mm, with 5 wavy-edged lobes, strong purplish red (61B) in bud, opening inside moderate purplish pink (N74D), with dark red (59A) spots on the dorsal lobes; outside strong reddish purple (72C); filaments white, anthers purple; style white, with a dark cherry red stigma. Calyx 1mm long, light green. Lvs elliptic, 125 × 60mm, rounded at base, pointed at apex, margins flat, glossy green above, hairless. Narrow bushy shrub, 2.4 × 2.0m in 20 years. Early June. Hardy to –30C.

Nomenclatural standard: colour print provided by the registrant (WSY0119762). Etymology: derived from the nursery name ("Babites") and a Latvian female Christian name ("Aiga").

## (r) 'Banana Cream'

cv. Elepidote rhododendron: 'Bisque' (s) × 'Capistrano'. **H:** J. Doppel (2008), **G:** J. Doppel (2011), **N:** J. Doppel (2014), **REG:** J. Doppel (2015).

Fls 12/domed truss, broadly funnel-shaped, 44mm long, with 5 rounded, wavy-edged lobes, buds deep yellowish pink (39B), becoming moderate yellowish pink (31D) just before opening, opening inside pale yellow-green (4D), with strong greenish yellow dorsal speckling, outside shading from dark greenish yellow (152D) at base to pale greenish yellow (2D) at margin; not scented. Calyx 3mm long, deep yellowish pink (39B). Lvs elliptic, 125 × 40mm, rounded at base, broadly acute at apex, margins flat, matt moderate olive green (137A) above, hairless. Dense shrub, 0.8 × 1.1m in 7 years. Midseason (May in eastern Pennsylvania, USA). Hardy to –23C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119838).

## (r) 'Barbara Lansdowne'

cv. Elepidote rhododendron: 'Meg Merrilees' (s) × unknown.

**G:** Marquis of Lansdowne (2000), **S:** Marquis of

Lansdowne (2000), N: Marquis of Lansdowne (2015), REG: I.T.Stokes (2015).

Fls 10/domed, slightly flattened lobes, funnel-shaped, 50 × 60mm, with 5 wavy-edged lobes, buds strong purplish red (63B), becoming paler towards the base, opening inside light purplish pink (55C), paling to pale pink (49D) towards the base, outside pale purplish pink (62D), darkening to strong purplish pink (62A) on the ribs; with a bold flare of deep red (60A) for half the length of the tube below the dorsal lobe; stamens hairy at base; ovary with short red glands which extend sparsely on to the base of the otherwise hairless style. Calyx 2–4mm long, teeth triangular, green, with a red glandular margin. Pedicels glandular. Lvs obovate, 70–100 × 30–40mm, cuneate to rounded at base, apex rounded and mucronate, margins flat, matt green above, hairless. Compact, somewhat rounded shrub, 1.5 × 4.0m in 25 years. Midseason (early/mid May). Hardy to –15C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119836). Etymology: named for the mother of the current Marquis of Lansdowne.

(r) **'Barb Blankenship'**

cv. *Elepidote rhododendron*: 'Independence Day' (s) × 'Peach Charm'.

H: J.Barlup (1996), G: J.Barlup (2002), N: J.Barlup (2012), REG: J.Barlup (2015).

Fls 19/ball truss, broadly funnel-shaped, 48 × 76mm, with 6 wavy-edged lobes, buds deep purplish pink (66D), opening inside strong yellow (153D), with a 6mm wide strong purplish pink (68D) marginal band on all lobes, the dorsal lobe with twin flares of strong purplish red (60B) extending 9mm from the base and surmounted by similarly coloured spots for 19mm; outside strong yellow (153D), with a 6mm wide deep purplish pink (68A) marginal band on all lobes, each lobe with a deep purplish pink (68A) central vein. Calyx 19mm long, strong yellow (153D), with strong purplish red (60B) spots and moderate purplish pink (68B) margins. Lvs elliptic, 127 × 51mm, rounded at base, broadly acute at apex, margins flat, slightly up-angled from the mid vein, dull and moderate olive green (147A) above, hairless. Shrub 2.0 × 2.4m in 18 years. Midseason (late May in Seattle, Washington, USA). Hardy to –18C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119689). Etymology: named for a friend of the registrant, a nurserywoman in Arlington, Washington, USA.

(a) **'Beautiful Beth'**

cv. *Evergreen azalea*: 'Ashley Ruth' × 'Komo-kulshan'.

H: J.F.Klimavicz (1999), G: J.F.Klimavicz (2001), N: J.F.Klimavicz (2015), REG: J.F.Klimavicz (2015).

Fls single, 2–3/single bud inflorescence, broadly funnel-shaped, 50mm wide, with 5 broadly rounded, very wavy-edged lobes, buds strong purplish pink (68B) at apex, shading to white at base, opening inside white in the centre with a strong purplish pink (63C & 68B) picotee band around all lobes, outside white at base with a broad, strong purplish pink (68B), picotee band around all lobes; filaments white;

style strong purplish pink (63C). Calyx 5mm long, strong yellow-green (144C). Lvs elliptic, 30 × 15mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Shrub 0.9 × 0.9m in 10 years, very floriferous. Midseason (early May, in Washington, D.C., USA). Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119848). Etymology: named for the mother-in-law of the hybridiser's daughter. Hybridiser's reference: 1K-99–4.

(r) **'Big Yellow'**

Syn. of 'Bob Furman's Big Yellow'.

(r) **'Bob Furman'**

cv. *Elepidote rhododendron*: believed to be a seedling from a Cecil Smith specimen of *yakushimanum*.

G: R.A.Furman (late 1990s), S: R.A.Furman (c. 2007), N: K.F.Humphries (2015), I: Van Veen Nursery (2013), REG: N.Beaudry (2015).

Fls 10–11/domed truss, campanulate, 43 × 48mm, with 5 wavy-edged lobes, deep purplish pink (54B) in bud, opening white (155D), with pale purplish pink (62D) in the centre of the lobes which fades to white (155D), and with moderate yellow-green (146D) to strong yellow-green (145A) spotting on the dorsal lobe; outside pale purplish pink (62D), shading to white (166D); not scented. Calyx insignificant. Lvs elliptic, 91 × 36mm, rounded at base, broadly acute at apex, margins downcurved, moderate olive green (147A) above, hairs below pale yellow (161C–D) at first, ageing to moderate orange-yellow (164B–165C); hairs above moderate orange-yellow (164B), persisting for 6–8 weeks. Dense shrub, 1.7 × 1.8m in 20 years. Midseason (June on Cape Cod, Massachusetts, USA). Hardy to –23C.

Nomenclatural standard: colour print of an electronic image provided by D.W. Hyatt at the time of registration (WSY0119834). Etymology: named for Bob Furman who grew the plant from seed and selected this clone. Originally circulated with the reference: Furman's R. yaku #1.

(r) **'Bob Furman's Big Yellow'**

cv. *Elepidote rhododendron*: 'Golden Star' × (Jalisco Group × *yakushimanum*) (s) × (Dido Group × ('Chlorops' × *lacteam*)) × 'Golden Star'.

H: R.A.Furman (1985), G: R.A.Furman (1992), N: R.A.Furman (1992), REG: J.Delano (2015).

Fls 12–14/domed truss, broadly funnel-shaped, 74 × 120mm, with 7 wavy-edged, emarginated lobes, waxy and with a heavy substance, moderate reddish orange (171A) fading to moderate orange-yellow (164B) in bud, opening inside light yellow (10B), with strong yellow-green (144B–C) spotting emanating from the throat and extending on to the three dorsal lobes; outside light orange-yellow (16C) to pale orange-yellow (16D); slightly scented. Calyx insignificant. Lvs elliptic, 140 × 53mm, rounded at base, obtuse at apex, margins slightly wavy, upangled from midvein, semiglossy and moderate olive green (147A) above, hairless. Shrub 2.7 × 2.7m in 29 years. Midseason (late May on Cape Cod, Massachusetts, USA). Hardy to –25C.

Syn: 'Big Yellow'.

Nomenclatural standard: colour print of an electronic image provided by Donna M. Delano (WSY0119833).

(r) 'Bradley Hayman'

cv. Elepidote rhododendron: 'The General' (s)  
× *fortunei*.

**H:** P.B.Moser (2007), **G:** P.B.Moser (2014),

**N:** P.B.Moser (2015), **REG:** P.B.Moser (2015).

Fls 6–10/domed truss, funnel-shaped, 51 × 76mm, with 6 rounded, wavy-edged lobes, strong purplish red (61B) in bud, opening inside very pale pink (73D), shading to light purplish pink (73C) at margins, with a small area of deep purplish red at the base; outside light purplish pink (73C). Calyx 2mm long, strong yellow-green (144C). Lvs elliptic, 127 × 60mm, rounded at base, obtuse at apex, margins flat, but lamina slightly concave, moderate olive green (146A) above, hairless. Shrub 1.1 × 0.8m in 8 years. Midseason (May in Philadelphia, USA). Hardy to –18C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119832).

Etymology: named for the deceased godson of the registrant.

(r) 'Carlostar'

cv. Elepidote rhododendron: 'Admiral Vanessa' (s)  
× 'Welse'.

**H:** K.-H.Hubbers (2000), **G:** K.-H.Hubbers (2007),

**N:** W.Wüstemeyer (2010), **I:** W.Wüstemeyer (2010),

**REG:** J. Westhoff (2015).

Fls c.15/domed truss, open funnel-shaped, 60 × 80mm, with 5–6, wavy and slightly decurved lobes, strong red (53C) in bud, opening inside light yellowish pink (27A) in the centre, shading out to strong to moderate pink (49A/B) near the margin, outside strong red to deep pink (51A/B) on the midribs, otherwise moderate to light pink (49B/C), shading up to light to pale yellowish pink (29C/D); with a large dark red (187A/B) blotch, solid at the base but densely spotted above, extending over a large part of the dorsal lobe, the markings just spreading to adjacent lobes. Calyx cut into irregular lobes, 8–23mm long, coloured like the inside of the corolla. Lvs oblong-lanceolate, 120 × 40mm, roundish at base, cuneate and with a small apiculus at apex, margins flat, matt green above, hairless. Shrub 1.2 × 1.1m in 10 years. Late May. Hardy to –23C.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0119830).

Etymology: named for the hybridiser.

(r) 'Dancefloor'

cv. Elepidote rhododendron: 'Fantastica' (s)  
× 'Double Eagle'.

**H:** H.Hesse (1999), **G:** H.Hesse (2004), **N:** H.Hesse, **I:** Baumschule H.Hachmann (2014), **REG:** H.Hesse (2015).

Fls 9–16/dense, rounded truss, broadly funnel-campanulate, 39–44 × 51–74mm, with 5 (-6) wavy-edged lobes, opening pale yellowish pink (29D) to light yellowish pink (36B & C), with a hint of strong pink (48C & D) at the margins, ageing to pale yellow (8D); outside strong red (51A), with parts pale

yellowish pink (29D) when first opening, and with the internal markings showing through, when fully open strong pink (48C & D) with pale yellowish pink (29D), ageing to pale greenish yellow (9D), with some strong pink (48C) parts and veins along the midvein; a mark of vivid red (45A) with black at the base of the dorsal lobe, with spots extending on to the middle of the dorsal lobe and below the other lobes (light red to strong red, 46A); filaments white, very short (one third the length of the style); style light green at base, shading to yellow-green at the apex. Calyx 1–2mm long, light green, with reddish edges. Lvs elliptic, 100–128 × 46–69mm, rounded at base, broadly acute at apex, margins flat, glossy green above, hairless. Rounded, upright shrub, 0.7 × 0.7m in 10 years. Early May to early June. Hardy to –20C. Nomenclatural standard: colour print provided by the registrant (WSY0119734).

(r) 'Dartmoor Evening Glow'

cv. Elepidote rhododendron: seedling from a selfed 'Spring Morning'.

**H:** B.N.Starling (1998), **G:** B.N.Starling (2001),

**N:** B.N.Starling, **I:** B.N.Starling (2012),

**REG:** B.N.Starling (2015).

Fls pendent, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 7mm, with 5 flat-edged lobes, deep salmon in bud, opening salmon inside and salmon rose outside; anthers dark brown; style pale green.

Calyx 3mm long, green. Pedicels with fine hairs. Lvs elliptic, 60 × 25mm, cuneate at base, acute at apex, margins flat, matt and glaucous above, hairy above, with a few bristles on the veins below; petioles sparsely hairy. Shrub with an open habit, 1.5 × 1.0m in 10 years, with hairy stems. May to June. Hardy to –15C. Previously listed as a *Menziesia* 'Evening Glow'.

Etymology: the Dartmoor prefix relates to the location of the registrant's garden in the UK, "Evening Glow" alludes to the colour of the flowers.

(r) 'Dartmoor Goldilocks'

cv. Elepidote rhododendron: derived from selfed 'Spring Morning'.

**H:** B.N.Starling (2009), **G:** B.N.Starling (2011),

**N:** B.N.Starling, **I:** B.N.Starling (2014),

**REG:** B.N.Starling (2015).

Fls pendent, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 8mm, with 5 flat-edged lobes, deep gold in bud, opening inside cream to gold; outside gold to deep gold, the lobes tinged with apricot; anthers dark brown; style cream. Calyx 3mm long, green. Pedicels with fine hairs. Lvs elliptic, 40 × 18mm, cuneate at base, acute at apex, margins flat, sparsely hairy. Shrub with a fairly open habit, 0.6 × 0.5m in 5 years. April. Hardy to at least –15C. Previously listed as *Menziesia* 'Goldilocks'. The Dartmoor prefix relates to the location of the registrant's garden in the UK, "Goldilocks" alludes to the flower colour.

(r) 'Dartmoor Moonlight'

cv. Elepidote rhododendron: 'Spring Morning'

× *menziesii*, glaucous form.

**H:** B.N.Starling (2005), **G:** B.N.Starling (2008),

N: B.N.Starling, I: B.N.Starling (2014),

**REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 7mm, with 5 flat-edged lobes, light brown to creamy green in bud, opening inside and out pale yellow; anthers dark brown; style creamy green. Calyx 2mm long, green. Pedicels with fine hairs. Lvs elliptic, 60 × 30mm, cuneate at base, acute at apex, margins wavy, matt and glaucous above, hairy above and with a few bristles on the veins below; petioles sparsely hairy. Shrub with an open habit, 1.5 × 1.0m in 10 years, with hairy stems. May to June. Hardy to at least –15C.

Previously listed as *Menziesia* ‘Moonlight’. The Dartmoor prefix relates to the location of the registrant’s garden in the UK, “Moonlight” alludes to the colour of the flowers.

(r) ‘**Dartmoor Shepherd’s Delight**’

cv. Elepidote rhododendron: seedling from selfed ‘Spring Morning’.

**H:** B.N.Starling (1991), **G:** B.N.Starling (1995),

**N:** B.N.Starling, **I:** B.N.Starling (2012),

**REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot tip, urceolate, with 5 flat-edged lobes, deep cream to rose pink in bud, opening inside cream, with rose pink lobes, outside rose pink, darkening towards the lobes; anthers dark brown; style cream; not scented. Calyx 2mm long, green. Pedicels with fine hairs. Lvs elliptic, 40 × 16mm, cuneate at base, acute at apex, margins flat, matt and glaucous above, hairy above and with a few bristles below; petioles sparsely hairy. Shrub with a fairly open, branching habit, 1.5 × 1.0m in 10 years, with hairy stems. May to June. Hardy to –15C.

Previously listed as *Menziesia* ‘Shepherd’s Delight’.

Etymology: the Dartmoor prefix relates to the location of the registrant’s garden in the UK, “Shepherd’s Morning” relates to the old saying ‘red sky at night, shepherd’s delight’, the flower colour being reminiscent of a sunset.

(r) ‘**Dartmoor Smoky Rose**’

cv. Elepidote rhododendron: seedling from selfed ‘Spring Morning’.

**H:** B.N.Starling (2002), **G:** B.N.Starling (2006),

**N:** B.N.Starling, **I:** B.N.Starling (2012),

**REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 6mm wide, buds smoky purple, with a yellow tip, opening smoky purple inside, outside smoky rose shading to rose at tips; anthers brown; style pale green. Calyx 3mm long, green. Pedicels with fine hairs. Lvs elliptic, 65 × 35mm, cuneate at base, acute at apex, matt and glaucous above, hairy above and with a few bristles on the veins below; petioles sparsely hairy. Open, upright shrub, 1.5 × 1.0m in 10 years, with hairy stems. May to June. Hardy to at least –15C.

Previously listed as *Menziesia* ‘Smoky Rose’. The Dartmoor prefix relates to the location of the registrant’s garden in the UK and “Smoky Rose” alludes to the colour of the flowers. The original

spelling “Smoky Rose” was an unintentional error and has, with the registrant’s approval, been amended to reflect the usual spelling of this adjective.

(r) ‘**Dartmoor Snowbells**’

cv. Elepidote rhododendron: seedling from selfed ‘Spring Morning’.

**H:** B.N.Starling (1995), **G:** B.N.Starling (2001),

**N:** B.N.Starling, **I:** B.N.Starling (2014),

**REG:** B.N.Starling (2015).

Fls pendent, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 6mm, with 5 flat-edged lobes, pale pink in bud, opening inside white, outside white with a faint pink rim; anthers pale fawn; style white to pale green. Calyx 3mm long, green. Pedicels with fine hairs. Lvs elliptic, 45 × 20mm, cuneate at base, acute at apex, margins flat, matt, silvery grey above, hairy above and with a few bristles on the veins below; petioles sparsely hairy. Slow-growing, compact shrub, 1.25 × 0.7m in 10 years, with hairy stems. May to June. Hardy to at least –15C.

Previously listed as *Menziesia* ‘Snowbells’.

Etymology: the Dartmoor prefix relates to the location of the registrant’s garden in the UK.

(r) ‘**Dartmoor Summer Skies**’

cv. Elepidote rhododendron: *multiflorum*, glaucous form × *menziesii*, glaucous form.

**H:** B.N.Starling (1986), **G:** B.N.Starling (1991),

**N:** B.N.Starling, **I:** B.N.Starling (1997),

**REG:** B.N.Starling (2015).

Fls pendent, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 8 × 5mm, with 5 flat-edged lobes, deep salmon pink in bud, opening inside apricot, the outside apricot shading to deep rose at the rim; anthers brown; style pale green. Calyx 2mm long, glaucous green. Pedicels with fine hairs. Lvs broadly elliptic, 50 × 35mm, cuneate at base, acute at apex, margins flat, matt and glaucous above, hairy above, with a few bristles on the veins below; petioles sparsely hairy. Shrub with an open, spreading habit, 1.5 × 1.5m in 10 years, with hairy stems. May to June. Hardy to at least –15C.

Previously listed as *Menziesia* ‘Summer Skies’.

Etymology: the Dartmoor prefix relates to the location of the registrant’s garden in the UK.

(r) ‘**Dartmoor Sunshine**’

cv. Elepidote rhododendron: ‘Dartmoor Moonlight’ × ‘Dartmoor Shepherd’s Delight’.

**H:** B.N.Starling (2005), **G:** B.N.Starling (2010),

**N:** B.N.Starling, **I:** B.N.Starling (2012),

**REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 6mm, with 5 flat-edged lobes, light brown to straw-coloured in bud, opening yellow inside, outside yellow with a pink rim; anthers dark brown; style pale green. Calyx 2mm long, green. Pedicels with fine hairs. Lvs elliptic, 60 × 25mm, cuneate at base, acute at apex, margins wavy, green above, sparsely hairy. Shrub with an open habit, 1.0 × 0.6m in 8 years. May to June. Hardy to at least –15C.

Previously listed as *Menziesia* ‘Sunshine’. The Dartmoor prefix relates to the location of the

registrant's garden in the UK, "Sunshine" alludes to the colour of the flowers.

(r) 'Dawn Debonair'

cv. *Elepidote rhododendron*: 'Winning One', open-pollinated.

G: J. Doppel (1998), S: J. Doppel (2014), N: J. Doppel (2014), REG: J. Doppel (2015).

Fls 16/domed truss, broadly funnel-shaped, 41 mm long, with 5 rounded, flat-edged lobes, with deep sinuses, deep purplish pink (N57C) in bud, opening inside light purplish pink (65B), with minor pale yellowish pink (29D) spotting on the dorsal lobe, outside deep purplish pink (64D); not scented. Calyx insignificant. Lvs elliptic, 125 × 50 mm, rounded at base, broadly acute at apex, margins flat, semiglossy and moderate olive green (137C) above, hairless. Shrub 1.5 × 1.2 m in 21 years. Midseason (May in eastern Pennsylvania, USA). Hardy to -26°C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119828).

Derived from open-pollinated seed collected by the registrant in 1994.

(r) 'Elsie Maria'

cv. *Elepidote rhododendron*: *dichroanthum* × unknown, open-pollinated.

H: R. Hahkala (1998), G: K. Theqvist (2002), N: K. Theqvist (2015), REG: K. Theqvist (2015).

Fls 14–15/rounded truss, funnel-shaped, 32–35 × 43–48 mm, with 5 wavy-edged lobes, strong purplish red (61B) in bud, opening inside white to pale purplish pink (65D), with deep purplish pink (N57C) on the edges which fades to moderate or light purplish pink (65A/B); outside white to pale purplish pink (65D), with vivid purplish red (N57B) stripes, deep purplish pink (N57C) on the edges which fades to moderate or light purplish pink (65A/B); pale yellow (8D) on the basal part of the dorsal lobe, with moderate yellow-green (146C) to moderate olive brown (199A) spotting; filaments white, anthers greyish purplish red (77D); style white, stigma strong red (46A). Calyx 1 mm long, light yellow-green (154D). Lvs oblanceolate to elliptic, 90–120 × 28–35 mm, cuneate at base, broadly acute at apex, margins sharply decurved, semi-glossy and moderate olive green (137A) above, hairless. Dense, low-growing shrub, 0.6 × 1.5 m in 15 years. Late season (late June to early July in Finland). Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119819).

Etyymology: named for the late mother of the registrant. The original *dichroanthum* hybrid was grown from open-pollinated seed on a plant of this species in Bremen Botanical Garden, Germany. This seedling was grown in the Turku Botanical Garden, in Ruissalo, Finland and open-pollinated seed from that plant gave rise to this new cultivar.

(r) 'Evening Breeze'

cv. *Elepidote rhododendron*: 'Bambi' × *proteoides* (s) × 'Plum Passion'.

H: J. Barlup (2005), G: J. Barlup (2009), N: J. Barlup (2014), REG: J. Barlup (2015).

Fls 13/domed truss, broadly funnel-shaped, 51 × 70 mm, with 5 wavy-edged lobes, buds deep red (60A), opening inside strong purplish pink (63C), blending to strong purplish red (63B) at margins, with strong purplish red (60C) midveins from base to tip, and with a moderate red (181A) dorsal flare extending 38 mm from the base; nectaries strong purplish red (60D); outside similarly coloured, but without the flare. Calyx 13 mm long, strong purplish pink (63C), with moderate red (181A) spotting on the tips. Lvs elliptic, 102 × 38 mm, rounded at base, broadly acute at apex, margins flat, moderate olive green (147A) above, hairless. Dense shrub, 0.6 × 0.9 m in 9 years. Midseason (May in Seattle, Washington, USA). Hardy to -18°C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119820).

(r) 'Farrah Parker'

cv. *Elepidote rhododendron*: 'Hearthside Beauty' (s) × *lukiangense* × 'August Moon'.

H: P. Smith (2001), G: P. Smith (2011), N: P. Smith (2013), REG: P. Smith (2015).

Fls 9–11/domed truss, open funnel-shaped, 51 × 51 mm, with 5–7 very wavy-edged lobes, strong purplish pink (55B) in bud, opening inside white, with light purplish pink (55C) mid veins and tips and a pale yellow (18C) blotch, with a few darker spots on the dorsal lobe, fading to all white; outside light purplish pink (55C), fading to white with light purplish pink (55C) mid veins. Calyx 2 mm long, green. Lvs obovate, 114 × 51 mm, cuneate at base, broadly acute at apex, margins flat, dark yellowish green (141B) above, hairless. Dense shrub, 0.8 × 0.9 m in 10 years. Early (February to March in northern California, USA), with a consistent and notable autumn flowering as well. Hardy to -8°C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119692).

Etyymology: named for a granddaughter of the registrant.

(r) 'Fiona Lansdowne'

cv. *Elepidote rhododendron*: Kewense Group (s) × 'Maxwell T. Masters'.

G: Marquis of Lansdowne (2000), S: Marquis of Lansdowne (2015), REG: I.T. Stokes (2015).

Fls 14–18/ tall, domed truss, open funnel-shaped, 60 × 80 mm, with 5–6 wavy-edged lobes, vivid purplish red (67B) in bud, opening inside strong reddish purple (72C), paling to light purplish pink (62C) on the inside of the ribs, outside vivid purplish red (66B) on the ribs and strong reddish purple (72C) between the ribs; with a strong purplish red (61B) dorsal flare, solid at the base, becoming spotted above; filaments white, hairy at base; ovary and base of style glandular. Calyx 1–4 mm long (uppermost lobe longest), green. Pedicels sparsely glandular. Lvs oblanceolate, 140–200 × 50–70 mm, rounded at base and apex, margins flat, matt green above, hairless. Large rounded shrub, 3.0 × 3.0 m in 20 years. Midseason (mid May in 2015). Hardy to -15°C.

Nomenclatural standard: colour print of an electronic



image provided by the registrant (WSY0119815).  
Etymology: named for the wife of the current Marquis of Lansdowne.

(r) **'Flaming Gold'**

cv. Elepidote rhododendron: sport from a plant of the Elizabeth Group.  
**S:** J. Goossens (1994), **N:** J. Goossens (1994),  
**I:** J. Goossens (1999), **REG:** J. Goossens (2015).  
Fls 6–8/open, flat truss, funnel-campanulate, 60 × 60mm, with 5 wavy-edged lobes, red throughout. Lvs elliptic, 70 × 25mm, margins flat, matt green above, with broad, irregular, variegated margins that are reddish brown at first, but the variegation ages to yellow, with an indumentum when young which appears to be lost with age. Shrub 1.2 × 2.0m in 10 years. Late April. Hardy to –18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119814).

(v) **'Flourish of Trumpets'**

cv. Lepidote rhododendron: 'Simbu Sunset' (s) × *suaveolens*.  
**H:** A. Rouse (2007), **G:** A. Rouse (2012), **N:** A. Rouse (2015), **REG:** A. Rouse (2015).  
Fls 16–20/hemispherical but slightly open-topped truss, trumpet-shaped, 48–50 × 30–34mm, with 5 very slightly wavy-edged lobes, light green in bud, opening white inside and out, unmarked; filaments white, anthers brownish orange; slightly scented. Calyx less than 1mm long, off white. Lvs broadly elliptic to oblanceolate, 25–55 × 10–30mm, broadly tapering to rounded at base, obtuse/rounded at apex (occasionally with an acuminate tip), blade flat, glossy green above, with tiny, sparse scales (not clearly visible to the naked eye). Stiff-branched shrub, 0.6 × 0.5m in 8 years. April to May (i.e. autumn in Australia).  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119811).

(r) **'Forrest Schoenberger'**

cv. Elepidote rhododendron: unknown (fls light yellow, with a red tinge) (s) × 'Kimberly' (Greer).  
**H:** P.B. Moser (2009), **G:** P.B. Moser (2015),  
**N:** P.B. Moser (2015), **REG:** P.B. Moser (2015).  
Fls 5–7/domed truss, open funnel-shaped, 44 × 70mm, with 7 emarginate, wavy-edged lobes, light purplish pink (63D) in bud, opening inside white (155C), with very pale pink (69B) margins, unmarked; outside white (155C), with very pale pink (69B) margins. Calyx 2mm long, strong yellow-green (144C). Lvs elliptic, 76 × 38mm, rounded at base, broadly acute at apex, margins upcurved, moderate yellow-green (137C) above, hairless. Dense shrub, 0.7 × 0.6m in 6 years. Midseason (May in Philadelphia, Pennsylvania, USA). Hardy to –18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119810).  
Etymology: named for the deceased nephew of the registrant.

(r) **GEOFF'S CELEBRATION**

Trade designation for 'GRAgff'.

(r) **'Gle028'**

cv. Elepidote rhododendron: 'Graziella' (Hachmann) × 'Loch Lomond'.  
**H:** P.A. & K.N.E. Cox (2003), **G:** P.A. & K.N.E. Cox, **N:** P.A. & K.N.E. Cox, **I:** Glendoick Gardens (2015), **REG:** Glendoick Gardens (2015).  
Fls to c.10/truss, funnel-shaped, with 5 hardly wavy lobes, flushed pale peachy pink to cream, like 'Percy Wiseman'; style and filaments whitish; anthers brown. Calyx very short. Pedicels reddish. Lvs narrow, dark green. Shrub with a dense, upright and compact habit, 0.8–1.35m tall in 10 years. Quite late flowering.  
Trade designation: GLENDICK SORBET.

(r) **'Gle029'**

cv. Elepidote rhododendron: 'Graziella' (Hachmann) × 'Loch Lomond'.  
**H:** P.A. & K.N.E. Cox (2003), **G:** P.A. & K.N.E. Cox, **N:** P.A. & K.N.E. Cox, **I:** Glendoick Gardens (2015), **REG:** Glendoick Gardens (2015).  
Fls strongly tinted yellowish pink in bud in upper part, suffused pale yellow towards base, opening pale yellow tinged pink, to almost white at margins and with stronger purplish pink staining right at the base inside; filaments white, anthers brown; style greenish white, stigma greenish. Calyx very short. Lvs very narrow, dark green. Shrub with a dense, compact habit, 0.8–1.34m tall in 10 years.  
Trade designation: GLENDICK SHERBET.

(r) **GLENDICK SHERBET**

Trade designation for 'Gle029'.

(r) **GLENDICK SORBET**

Trade designation for 'Gle028'.

(r) **'Golden Radiance'**

cv. Elepidote rhododendron: 'Scintillation' × 'Phipps Yellow' (s) × 'Bud's Yellow'.  
**H:** H.C. Gehrlich (2004), **G:** H.C. Gehrlich (2011), **N:** H.C. Gehrlich (2014), **REG:** H.C. Gehrlich (2015).  
Fls 14/ball truss, open funnel-shaped, 51 × 64mm, with 7 wavy-edged lobes, buds light greenish yellow (4B), opening light greenish yellow (4B) throughout. Calyx insignificant. Lvs oblong, 127 × 48mm, rounded at base, obtuse at apex, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub 1.7 × 1.2m in 10 years. Midseason (May on Long Island, New York, USA). Hardy to –18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119809).

(r) **'GRAgff'**

cv. Elepidote rhododendron: 'Jeanette Clarke' (s) × *arboresum*.  
**H:** L. Begg (2003), **G:** J.C. Gray (2010), **N:** J.C. Gray (2014), **I:** Brindabella Country Gardens (2015), **REG:** J.C. Gray (2015).  
Fls 12/ball truss, campanulate, 55 × 50mm, with 5 very wavy-edged lobes, deep yellowish pink (46D) in bud, opening deep purplish pink (55A) throughout, with crimson markings on the dorsal lobe. Lvs elliptic, 110 × 40mm, margins upcurved, glossy

green above, hairless. Shrub 1.2 × 1.0m in 10 years. September (Australia). Very heat tolerant. Trade designation: GEOFF'S CELEBRATION. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119694).

(r) 'Helen White'

cv. Elepidote rhododendron: parentage unknown. **H:** G.P.White (2003), **G:** G.P.White, **N:** G.P.White, **REG:** G.P.White (2015). Fls 16/low domed truss, funnel-shaped, 45 × 35mm, with 5 wavy-edged lobes, dark purplish pink in bud, opening inside pale purplish pink, shading much paler in the centre of the lobes and in the throat, with conspicuous, orange, dorsal spotting; stamens white; style white, stigma very pale yellow. Calyx short. Lvs elliptic, 140 × 40mm, cuneate at base, broadly acute at apex, margins decurved, glossy green above, with white hairs when young which age to pale green. Shrub 1.2 × 1.0m in 10 years. Late May. Hardy to -20C. Nomenclatural standard: colour print provided by the registrant (WSY0119660). Etymology: named for one of the twin daughters of the registrant.

(r) 'Holly's Faith'

cv. Elepidote rhododendron: 'Christina Dee' (s) × 'My Jane'. **H:** J.Doppel (2008), **G:** J.Doppel (2011), **N:** J.Doppel (2013), **REG:** J.Doppel (2015). Fls 14/conical truss, broadly funnel-shaped, with 5 broadly acute, wavy-edged lobes, pale yellow-green (4D) in bud, opening inside light greenish yellow (8C), with a moderate orange (167A) speckled blotch on the dorsal lobe, which ages to green; outside pale yellow-green (4D). Calyx insignificant. Lvs elliptic, 127 × 38mm, rounded at base, broadly acute at apex, margins downcurved, semiglossy and moderate yellow-green (137C) above, hairless. Dense shrub, 0.6 × 0.8m in 7 years. Midseason (early May in eastern Pennsylvania, USA). Hardy to -23C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119807). Etymology: named for the wife of the registrant.

(r) 'Ilgonis'

cv. Elepidote rhododendron: 'Purple Splendour' (s) × 'Penenka'. **H:** R.Kondratovičs (1987), **G:** R.Kondratovičs (1993), **N:** R.Kondratovičs (2005), **REG:** Rhododendron Nursery Babite, University of Latvia (2015). Fls 14/strongly domed truss, open funnel-shaped, 75–80mm wide, with 5–6 wavy-edged lobes, buds strong reddish purple (N78B), opening inside light purple (N78D), with dark purplish red (N79A) spots on the dorsal lobes, outside strong reddish purple (N78B); filaments light purple, anthers white; style white, with a dark purple stigma. Calyx 5mm long, light purple. Lvs elliptic, 140 × 65mm, rounded at base, pointed at apex, margins wavy, glossy green above, hairless. Broad, bushy shrub, 1.7 × 2.2m in 25 years. Late May to early June. Hardy to -25C. Nomenclatural standard: colour print provided by the registrant (WSY0119763). Etymology: the epithet is a male Latvian Christian name.

(a) 'Illusia'

cv. Deciduous azalea: 'Bouquet de Flore' (s) × unknown yellow-flowered hybrid from Arboretum Mustila (probably involving *luteum*). **H:** A.Väinölä (1988), **G:** University of Helsinki and City of Helsinki, **N:** A.Väinölä (2009), **I:** MTT Agrifood Research Finland (2010), **REG:** University of Helsinki (2015). Fls 10–13/flat truss, very narrowly funnel-shaped, 45–50 × 24–45mm, with 5 wavy-edged lobes, buds pale yellow (18D, 158B), opening white (155D) inside and out, dorsal lobe strong orange-yellow (17A); scented. Calyx 1–2mm long, light green. Lvs elliptic, 57–69 × 20–24mm, cuneate at base, acute at apex, margins decurved, matt green above, hairy, with good reddish autumn leaf colour. Shrub 1.3 × 1.4m in 20 years. Mid June in Finland. Hardy. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119808). Etymology: named for a fairy in a Finnish fairytale by Y.Kokko (1944). First published in *Sorbifolia* 43(1):34–38 (2012).

(r) 'Jessica Gustavson'

cv. Elepidote rhododendron: 'Violet Mist' (s) × 'Plum Passion'. **H:** J.Barlup (1995), **G:** J.Barlup (2000), **N:** J.Gustavson (2015), **REG:** J.Barlup (2015). Fls 15/ball truss, broadly funnel-shaped, 47 × 67mm, with 5 deep, wavy-edged lobes, vivid purple (81B) in bud, opening inside light purple (84C), shading to strong purple (84A) at all margins, with a large greyish red (182B) speckled blotch superimposed over dark greenish yellow (152C) on the dorsal lobe; outside light purple (84C) at base, shading to strong purple (84A) at margins; filaments white; style reddish; not scented. Calyx insignificant. Lvs elliptic, 102 × 44mm, broadly acute at apex, rounded at base, margins flat, matt and moderate olive green (147A) above, hairless. Open shrub, 0.9 × 0.9m in 20 years. Midseason (early May in Seattle, Washington, USA). Hardy to -15C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119846). Etymology: nominant purchased the right to name this cultivar at the American Rhododendron Society convention in 2015.

(a) 'Joe Klimavicz'

cv. Evergreen azalea: 'Haru-no-sono' (s) × 'Dorothy Clark'. **H:** Robert Stewart (1998), **G:** Robert Stewart (2001), **N:** Robert Stewart (2013), **REG:** Robert Stewart (2015). Fls 2–3/cluster, broadly funnel-shaped, 51 × 76–90mm, with 5–6 rounded, wavy-edged and overlapping lobes, buds deep pink (48B) blending with strong pink (48C and 48D), with random areas of white, opening inside deep pink (48B) blending with strong pink (48C and 48D), with random areas of white and with strong yellow-green (145A) spotting on a white ground in dorsal part of flower; outside deep pink (48B) blending with strong pink (48C and 48D), with random areas of white; style

and filaments red; not scented. Calyx 6mm long, strong yellow-green (143C). Lvs elliptic, 51 × 25mm, rounded at base, broadly acute at apex, margins flat, moderate yellow-green (137C in spring, 147A in summer), hairless. Open shrub, 0.8 × 1.1m in 10 years. Midseason (mid May in Washington, D.C., USA). Hardy to -18C.

Nomenclatural standard: colour print of an electronic image provided by C. Beck (WSY0119805).

Etymology: named for a fellow hybridiser and friend of the registrant.

(r) **'Katy White'**

cv. Elepidote rhododendron: parentage unknown.

**H:** G.P.White (2003), **G:** G.P.White, **N:** G.P.White, **REG:** G.P.White (2015).

Fls 14/low domed truss, funnel-shaped, 45 × 35mm, with 5 wavy-edged lobes, deep purplish pink in bud, opening inside mid purplish pink, becoming much paler into the throat, with conspicuous, orange, dorsal spotting; stamens white; style white, stigma very pale yellow. Calyx short. Lvs elliptic, 140 × 40mm, cuneate at base, broadly acute at apex, margins decurved, glossy green above, with bronze hairs. Shrub 1.0 × 1.0m in 10 years. Mid May. Hardy to -20C.

Nomenclatural standard: colour print provided by the registrant (WSY0119661). Etymology: named for one of the twin daughters of the registrant.

(r) **'Ken Gibson'**

cv. Elepidote rhododendron: 'Lemon Embers' (s) × Fortune Group × ('Berg's Yellow' × *proteoides*).

**H:** J.Barlup (2003), **G:** J.Barlup (2013), **N:** J.Barlup (2014), **REG:** J.Barlup (2015).

Fls 19/ball truss, broadly funnel-shaped, 51 × 76mm, with 5 wavy-edged lobes, buds light greenish yellow (4C), opening inside light greenish yellow (4C) in centre, blending to pale yellow-green (4D), with a 13mm long strong red (53C) flare on the dorsal lobe; outside light greenish yellow (4C). Calyx 19mm long, light greenish yellow (4C), with a 6mm long strong red (53C) flare. Lvs elliptic, 121 × 60mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.4 × 1.4m in 12 years. Midseason (mid May in Seattle, Washington, USA). Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a noted Canadian rhododendron collector.

(r) **'Kristian's Pink'**

cv. Elepidote rhododendron: *brachycarpum* (as subsp. *tigerstedtii*) (s) × probably 'Roseum Elegans'.

**H:** Arboretum Mustila (1998), **G:** K.Theqvist (2003), **N:** K.Theqvist (2008), **REG:** K.Theqvist (2015).

Fls 12-14/rounded truss, funnel-shaped, 35-40 × 50-55mm, with 5 wavy-edged lobes, deep purplish red (61A) in bud, opening inside deep purplish pink (N66C), outside deep purplish pink (N66C); with a small white centre, and with deep greenish yellow (153A) dorsal spotting; filaments light red-purple, anthers white; style and stigma red. Calyx 1mm long, yellow-green. Lvs elliptic, 100-125 × 45-65mm, margins decurved, broadly acute at apex, rounded at

base, matt moderate olive green (137A) above, with a white tomentum on the upper surface, changing to light brown in autumn and disappearing with age; small brown hairs on the midvein below; petioles with small white hairs. Shrub 0.6 × 1.5m in 15 years. Midseason (mid June in Finland). Diploid (as shown by flow cytometry). Resembles 'Roseum Elegans' except for smaller leaves, lower growth habit and superior hardiness.

Syn: 'Roseum Mustila'.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119796).

Etymology: named for the registrant and the colour of the flowers.

(r) **'Lemon Lollipop'**

cv. Elepidote rhododendron: 'Austin's Gold' (s)

× 'Capistrano'.

**H:** J.Doppel (2009), **G:** J.Doppel (2012), **N:** J.Doppel (2013), **REG:** J.Doppel (2015).

Fls 12/domed truss, funnel-shaped, 44mm long, with 5 rounded, wavy-edged lobes, buds moderate orange (167B), becoming brilliant greenish yellow (151D) just before opening, opening inside light greenish yellow (7D), with a strong red (53C), star-shaped, basal flare; outside light greenish yellow (7D). Calyx 6mm long, strong yellow-green (141D) at base, brilliant yellow-green (149A) at margin. Lvs elliptic, rounded at base, broadly acute at apex, margins flat, matt and dark green (136A) above, hairless. Dense shrub, 0.5 × 0.7m in 6 years. Midseason (early May in eastern Pennsylvania, USA). Hardy to -23C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119795).

(r) **'Lemon Sun'**

cv. Elepidote rhododendron: 'Recital' (s) × Fortune Group × ('Berg's Yellow' × *proteoides*).

**H:** J.Barlup (2003), **G:** J.Barlup (2008), **N:** J.Barlup (2014), **REG:** J.Barlup (2015).

Fls 21/ball truss, funnel-campanulate, 51 × 70mm, with 6 wavy-edged lobes, buds moderate red (47A), opening inside pale yellow-green (4D), with light greenish yellow (4C) ridges and lobe edges, and with a strong red (53D) dorsal flare extending 9mm from the base; outside similarly coloured, but lacking the flare; contrasting orange-red anthers. Calyx 25mm long, pale yellow-green (4D), with a strong red (53D) flare. Lvs elliptic, 157 × 54mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.7m in 10 years. Midseason (late April in Seattle, Washington, USA). Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119794).

(r) **'Len's Pacific Foam'**

cv. Elepidote rhododendron: 'Golden Anniversary' (s) × 'Faggetter's Favourite'.

**H:** L.Charvet (1990s), **G:** R. & S.Trujillo (2000), **N:** Forest Lane Nursery (2014), **I:** Forest Lane Nursery (2014), **REG:** Forest Lane Nursery (2015). Fls 24/conical truss, broadly funnel-shaped, 60 × 97mm, with 7 flat-edged lobes, light pink (49C) in

bud, opening inside white (155A), with moderate red (180A) marking on the dorsal lobe and light pink (49C) at lobe tips which quickly fades; outside white (155A); slightly scented. Calyx insignificant. Lvs elliptic, 178 × 55mm, cuneate at base, broadly acute at apex, margins flat, moderate olive green (137A) above, hairless. Shrub 1.7 × 1.2m in 15 years. Midseason (April to May in coastal northern California, USA). Hardy to -7C.  
Etymology: named in part after the hybridiser, Len Charvet.

(r) 'Loey Jane'

cv. Elepidote rhododendron: 'Percy Wiseman' (s) × Fortune Group × ('Berg's Yellow' × *proteoides*).  
H: J. Barlup (2003), G: J. Barlup (2008), N: J. Barlup (2015), REG: J. Barlup (2015).

Fls 21/ball truss, broadly funnel-shaped, 44 × 70mm, with 7 wavy-edged lobes, strong red (53C) in bud, opening inside pale yellow-green (4D), blending on the dorsal lobes to light pink (49C) 13mm from the margin (most evident on the central dorsal lobe), with two 19mm long flares of strong red (53B) spots in the dorsal area; outside pale yellow-green (4D), blending to light greenish yellow (4C) at base and to light pink (49C) at the margins; not scented. Calyx 13mm long, pale yellow-green (4D). Lvs elliptic, 133 × 54mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.5 × 0.6m in 5 years. Midseason (late April in Seattle, Washington, USA). Hardy to -18C.  
Nomenclatural standard; colour print of an electronic image provided by the registrant (WSY0119847).  
Etymology: named for Lois Jane Blackmore, of Victoria, British Columbia, Canada.

(r) 'Lori's Love'

cv. Elepidote rhododendron: 'Hotei' × *yakushimanum* (s) × 'Recital'.

H: J. Barlup (1997), G: J. Barlup (2003), N: J. Barlup (2014), REG: J. Barlup (2015).

Fls 19/ball truss, broadly funnel-shaped, 44 × 64mm, with 5 wavy-edged lobes, buds deep pink (52B), opening inside pale yellow-green (4D), blended to light greenish yellow (4C) at the throat, and with twin, deep pink, dorsal flares at the base; outside pale yellow-green (4D), blending to light greenish yellow (4C) at the base, with prominent light yellowish pink (36B) areas along each midvein. Calyx 19mm long, pale yellow-green (4D), blending to light greenish yellow (4C) at base. Lvs elliptic, 90 × 44mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.8m in 6 years. Midseason (mid May in Seattle, Washington, USA). Hardy to -18C.

Nomenclatural standard; colour print of an electronic image provided by the registrant (WSY0119793).

(r) 'Mammadaba'

cv. Elepidote rhododendron: 'Kristine' (Kondratovičs) (s) × unknown.

H: R. Kondratovičs (1995), G: R. Kondratovičs

(2001), N: R. Kondratovičs (2015),

REG: Rhododendron Nursery Babites, University of Latvia (2015).

Fls 17/strongly domed truss, open funnel-shaped, 100 × 90mm, with 5 wavy-edged lobes, light purple (77C) in bud, gradually fading to white inside and out after opening, with strong greenish yellow (151A) spots on the dorsal lobe internally; filaments white, anthers white; style light green, with a pink stigma. Calyx 2mm long, light pink. Lvs elliptic, 150 × 70mm, rounded at base, pointed at apex, margins wavy, matt green above, hairless. Broad, bushy shrub, 2.4 × 3.0m in 20 years. Late May to early June. Hardy to -25C.  
Nomenclatural standard; colour print provided by the registrant (WSY0119764).  
Etymology: epithet means "Mother Nature" in Latvian.

(r) 'Mara'

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' (s) × unknown.

H: R. Kondratovičs (1963), G: R. Kondratovičs (1969), N: R. Kondratovičs (1969), REG: Rhododendron Nursery Babite, University of Latvia (2015).

Fls 14/slightly domed truss, funnel-shaped, 60mm wide, with 5 slightly wavy-edged lobes, strong reddish purple (72C) in bud, opening inside deep purplish pink (73A) at the margins, with slightly darker tips to the dorsal lobes, the centre of the flower paling to very pale purple (69B), and with strong greenish yellow (151A) spots on the dorsal lobes; outside deep purplish pink (73A); filaments light pink, anthers light purple; style light pink, with a yellow-green stigma. Calyx 2mm long, light green. Lvs elliptic, 150 × 50mm, pointed at base and apex, margins flat to slightly wavy, matt green above, hairless. Broad, bushy shrub, 2.4 × 2.4m in 17 years. Late May to early June. Hardy to -30C.  
Nomenclatural standard; colour print provided by the registrant (WSY0119765).  
Etymology: epithet is a female Latvian Christian name.

(a) 'Midsummer Coral'

cv. Deciduous azalea: 'Denny's Rose' × *viscosum* (selection no. 3).

H: E. G. Millais (2000), G: D. Millais (2004),

N: D. Millais (2015), I: D. Millais (2015),

REG: D. Millais (2015).

Fls 7-8/rounded truss, tubular funnel-shaped, 40-45 × 40-50mm, with 5 flat-edged lobes, buds deep yellowish pink (39B) at base, shading to red/pink (47) at the apex, opening inside strong pink (48D), with a darker line (deep pink, 48B) down the centre of each lobe, dorsal lobe with a large strong yellowish pink (32C) flare, outside deep pink (48A) at base, shading to strong pink (48C) on the lobes; scented. Calyx 2mm long, light green. Lvs elliptic, 45-75 × 20-30mm, margins decurved, glossy, light green, slowly turning glaucous blue, hairy. Shrub 1.5 × 1.2m in 10 years. Late June. Hardy to -20C.

Nomenclatural standard; colour print of an electronic image provided by the registrant (WSY0119792).

Etymology: the epithet refers to the time of flowering and flower colour. First published: press release from Millais Nurseries, for RHS Hampton Court Palace Flower Show, 2015.

(a) **'Midsummer Girl'**

cv. Deciduous azalea: 'Denny's Rose' (s) × *viscosum* (selection no. 1).

**H:** E.G.Millais (2000), **G:** D.Millais (2004),

**N:** D.Millais (2015), **I:** D.Millais (2015),

**REG:** D.Millais (2015).

Fls 9–10/domed truss, tubular funnel-shaped, 45–50 × 35–40mm, with 5 flat-edged lobes, deep purplish pink (54B) in bud, opening inside strong purplish pink (55B), changing to light purplish pink (55C), and with a strong orange-yellow (24B) flare on the much broader, dorsal lobe; outside strong purplish pink (55B), with darker midribs (deep purplish pink, 55A); scented. Calyx 5–6mm long, light green. Lvs elliptic, 55–65 × 18–22mm, margins decurved, hairy, light green, with a bronzy sheen in summer, and glaucous undersides. Shrub 1.5 × 1.2m in 10 years. Very late June to early July. Hardy to –20C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119791). Etymology: the epithets refer in part to the flowering time. First published in press release from Millais Nurseries for RHS Hampton Court Palace Flower Show, 2015.

(a) **'Midsummer Moon'**

cv. Deciduous azalea: 'Denny's Rose' (s) × *viscosum* (selection no. 1).

**H:** E.G.Millais (2000), **G:** D.Millais (2004),

**N:** D.Millais (2015), **I:** D.Millais (2015),

**REG:** D.Millais (2015).

Fls 9–10/domed truss, tubular funnel-shaped, 45–50 × 45–50mm, with 5 wavy-edged lobes, brilliant yellow (8A) in bud, opening inside light greenish yellow (8C), shading to pale yellow (8D), then fading to yellowish white (158D), with light greenish yellow (8C) markings on the dorsal lobe; outside pale yellow (8D), with deeper midribs (light greenish yellow, 8B); sweetly scented. Calyx 5mm long, light green. Lvs elliptic, 60–80 × 20–25mm, margins decurved, hairy, pale glossy green above in spring, glaucous blue below. Shrub 1.5 × 1.2m in 10 years. Late June to early July. Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119790).

Etymology: the epithet refers to the period of flowering and the colour of the flowers. First published in press release from Millais Nurseries for RHS Hampton Court Palace Flower Show, 2015.

(a) **'Midsummer Rose'**

cv. Deciduous azalea: 'Denny's Rose' (s) × *viscosum* (selection no. 3).

**H:** E.G.Millais (2000), **N:** D.Millais (2015),

**I:** D.Millais (2015), **REG:** D.Millais (2015).

Fls 8–13/rounded truss, tubular funnel-shaped, 40–45 × 40–50mm, with 5 wavy-edged lobes, strong purplish red (54A) in bud, opening strong purplish pink (68B) at edges, paler in the centre (light purplish pink, 68D), fading to pale purplish pink (65D), and with a faint vivid yellow (12A) flare on the dorsal lobe; outside strong to light purplish pink (68B–D); scented. Calyx 1–2mm long, pale green. Lvs elliptic, 55–80 × 20–25mm, margins decurved,

hairy, dark, glossy green. Shrub 1.5 × 1.2m in 10 years. Late June. Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119789).

Etymology: the epithet refers to the period of flowering and the colour of the flowers. First published in press release from Millais Nurseries for RHS Hampton Court Palace Flower Show, 2015.

(a) **'Midsummer Star'**

cv. Deciduous azalea: 'Denny's Rose' (s) × *viscosum* f. *rhodanthum*.

**H:** E.G.Millais (2000), **G:** D.Millais (2004),

**N:** D.Millais (2015), **I:** D.Millais (2015),

**REG:** D.Millais (2015).

Fls 9–11/domed truss, narrowly tubular funnel-shaped, 45–50 × 45–50mm, with 5 wavy-edged lobes, strong purplish red (63B) in bud, opening moderate purplish pink (62B), with slight yellow-green markings on the dorsal lobe, outside moderate purplish pink (62B), with a deeper pink tube (strong purplish pink, 62A); scented. Calyx 1–2mm long, light green. Lvs elliptic, 60–70 × 17–22mm, margins decurved, glossy green above, hairy. Shrub 1.5 × 1.2m in 10 years. Late June. Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119787).

Etymology: the epithet refers to the period of flowering and the shape of the 'face' of the flowers. First published in press release from Millais Nurseries for RHS Hampton Court Palace Flower Show, 2015.

(a) **'Midsummer Wedding'**

cv. Deciduous azalea: *arborescens* (s) × 'Denny's Rose'.

**H:** E.G.Millais (2000), **G:** D.Millais (2004),

**N:** D.Millais (2015), **I:** D.Millais (2015),

**REG:** D.Millais (2015).

Fls 8–10/rounded truss, long tubular funnel-shaped, 40–45 × 50mm, with 5 wavy-edged lobes, light yellow-green (150D) in bud, opening inside white (155D), with a split, yellow, dorsal flare, which starts vivid yellow (13A) and fades to pale greenish yellow (13D), outside white (155D); scented. Calyx 2–3mm long, pale green. Lvs elliptic, 60–90 × 23–36mm, margins decurved, matt and slightly glaucous above, glaucous blue below, hairy. New growth reddish. Shrub 1.5 × 1–2m in 10 years. Late June to early July. Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119788).

Etymology: the epithet refers to the period of flowering and the colour of the flowers (as in a white wedding). First published in press release from Millais Nurseries for RHS Hampton Court Palace Flower Show, 2015.

(r) **'Mondflimmern'**

cv. Elepidote rhododendron: 'Peste's Fire Light' (s) × 'Mooreule'.

**H:** H.Hesse (1999), **G:** H.Hesse (2005), **N:** H.Hesse (2009), **REG:** H.Hesse (2015).

Fls 13–18/slightly loose, rounded truss, open funnel-shaped, 45–58 × 72–96mm, with 5–7 wavy-edged lobes, opening inside pale yellow (19C & 20D), with

the base light yellow (12C), ageing to pale yellow (20D) and pale orange-yellow (23D), with a light yellow (12C) base; outside moderate red to deep yellowish pink (47A–C) and light yellowish pink (27A) when first starting to open, eventually opening pale yellow (20D) (with internal markings showing through), ageing to pale orange-yellow (23D); conspicuous dorsal markings of dark red (87A) which extend on to the adjacent lobes, with additional vivid red (46B) basal marks; filaments light yellow; style light yellow at base, shading to reddish at apex. Calyx 1–30mm long, irregular and petaloid, pale yellow (20D) and light yellow (12C), with dark red (187A) spots and vivid red (46B) basal marks. Lvs elliptic to broadly ovate, 117–139 × 51–59mm, rounded at base, acute at apex, matt green above, hairless. Broadly erect shrub, 1.2 × 1.0m in 10 years. Mid May to mid June. Hardy to –21C. Nomenclatural standard: colour print provided by the registrant (WSY0119737).

(r) **'Mrs Carolina Lago'**

cv. Elepidote rhododendron: 'Maverick' (s)  
× 'Crimson Constellation'.

**H:** J. Barlup (2005), **G:** J. Barlup (2010), **N:** J. Barlup (2014), **REG:** J. Barlup (2015).

Fls 17/ball truss, broadly funnel-shaped, 44 × 76mm, with 5 very wavy-edged lobes, buds strong red (53D), opening white (155D), blending through pale purplish pink (55D) to deep pink (55A) at margins, with strong red (53C) dorsal spotting and strong red (53C) nectaries; outside white (155D), blending through pale purplish pink (55D) to deep pink (55A) at margins. Calyx 25mm long, white (155D), with strong red (53C) spotting and deep pink (55A) margins. Lvs elliptic, 108 × 22mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 8 years. Midseason (early May in Seattle, Washington, USA). Hardy to –18C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119785). Etymology: named for a friend of the registrant from Alexandria, Virginia, USA.

(r) **'Noyo Mist'**

cv. Lepidote rhododendron: 'Butterhorn' (s) × 'Ila Mae Allenby'.

**H:** P. Johnson (2004), **G:** P. Johnson (2009), **N:** P. Johnson (2015), **REG:** P. Johnson (2015).

Fls 5/lax truss, funnel-shaped, 66 × 67mm, with 5 wavy-edged lobes, buds moderate yellow (160A), opening white (155A), with yellow in the throat; outside brilliant yellow-green (154C), fading to white (155A); moderately scented. Calyx 22mm long, strong yellow-green (145A). Lvs obovate, 50 × 28mm, rounded at base, obtuse at apex, margins downcurved, moderate olive green (147A) above. Shrub 0.9 × 0.9m in 11 years. Midseason (April to May in coastal northern California, USA). Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119786).

(v) **'Olivia Manson'**

cv. Lepidote rhododendron: 'Simbu Sunset' (s)  
× *luraluense*.

**H:** A. Rouse (2002), **G:** A. Rouse (2012), **N:** A. Rouse (2015), **REG:** A. Rouse (2015).

Fls 5–8/open truss (the flowers sometimes all pointing in the same direction), tubular funnel-shaped, 60–62 × 60–62mm, with 5 wavy-edged lobes, light green in bud, opening inside cream, with the slightest hint of lime green, flushed to light pink (49C); outside cream, with a slight hint of lime green on the tube, flushed to light pink on edges of the lobes; unmarked; filaments cream, flushed to light pink (49C), anthers dark brown; not scented. Calyx less than 1mm long, very pale lime green. Lvs broadly elliptic to oblanceolate, 50–70 × 18–40mm, broadly tapering at base, obtuse to acute, or occasionally rounded at apex, tip often reflexed, glossy green, with tiny, sparse scales (not clearly visible to the naked eye). Scrub 1.5m × 0.4m in 13 years. April to May (i.e. autumn-flowering in Australia).

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119777).

Etymology: named for the niece of the registrant on her 21st birthday.

(r) **'Onica'**

cv. Elepidote rhododendron: 'Sumner's Orange' (s)  
× 'Double Eagle' (Korth).

**H:** P. Smith (2001), **G:** P. Smith (2011), **N:** P. Smith (2013), **REG:** P. Smith (2015).

Fls 16–18/domed truss, open funnel-shaped, 51 × 76mm, with 5–7 wavy-edged lobes, strong purplish red (54A) in bud, opening from light yellowish pink (36B) at margin and blending through pale orange-yellow (16D) to vivid yellow (16A), with a 25mm wide throat of moderate red (47A); outside primarily light yellowish pink (36A), with light orange-yellow (16B) at the base. Calyx 3mm long, green. Lvs oblong, 114 × 38mm, rounded at base, broadly acute at apex, margins flat, dull and dark yellowish green (139A) above, hairless. Shrub 1.2 × 0.9m in 10 years. Midseason (March to April in northern California, USA). Hardy to –8C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119706).

Etymology: named for a daughter of the registrant.

(a) **'Onnimanni'**

cv. Deciduous azalea: 'Nancy Waterer' × unknown yellow-flowered hybrid (possible involving *luteum*).

**H:** A. Väinölä (1988), **G:** University of Helsinki and City of Helsinki, **N:** A. Väinölä (2010), **I:** MTT Agrifood Research Finland (2012), **REG:** University of Helsinki (2015).

Fls 16–17/low-domed truss, narrow funnel-shaped, 45 × 50–55mm, with 5 flat to wavy-edged lobes, buds vivid yellow to strong orange-yellow (15A to 17A–B) and vivid to strong reddish orange (43b, 42C), opening inside brilliant yellow to light yellow (10A, 11A–B and 14C), outside light to brilliant yellow (12C, 13C) and light orange-yellow and vivid reddish orange-yellow (16B–C, 43B); dorsal lobe covered by markings of strong to vivid orange-yellow

(17A, 21A and 23A); style and stamens yellow. Calyx 1–2mm long, light green. Lvs elliptic, 103 × 30mm, cuneate at base, acute at apex, margins flat to wavy; hairy. Shrub 1.3–1.45 × 1.5–1.65m in 15 years. Mid June. Hardy.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119778).  
Etymology: named after a character in a Finnish nursery rhyme (a green boy in the woods). First published in *Sorbifolia* 43(1):34–38 (2012).

(r) **‘Orange Taffy’**

cv. *Elepidote rhododendron*: ‘Southern Skies’ (s) × ‘Percy Wiseman’ × ‘Orange Marmalade’.

**H:** D.S.Wallace (1999), **G:** D.S.Wallace (2004), **N:** D.S.Wallace (2015), **I:** Singing Tree Gardens & Nursery (2015), **REG:** D.S.Wallace and Singing Tree Gardens & Nursery (2015).

Fls 13/domed truss, funnel-shaped, 64 × 76mm, the corolla with 5 very wavy-edged lobes, strong purplish red (60D) in bud, opening inside light greenish yellow (6D), with strong purplish pink (62A) at margins and deep purplish red (61A) at the base which shades to strong purplish red (63A) 19mm from the base; outside pale yellowish pink (27C), shading to moderate purplish pink (62B) at 13mm from the margin; not scented. Calyx 44mm long, pale yellowish pink (27C), shading to moderate purplish pink (62B). Lvs elliptic, 140 × 38mm, rounded at base, broadly acute at apex, margins downcurved, the blade strongly convex, semiglossy and moderate bluish green (127A) above, hairless. Shrub 1.2 × 1.5m in 10 years. Midseason (April in coastal northern California, USA). Hardy to –18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119779).  
Breeder’s ref.: W9930.

(a) **‘Perfectly Pauline’**

cv. Deciduous azalea: parentage unknown.

**H:** W.G.T.Hyde (1970s), **G:** W.G.T.Hyde (1970s), **N:** Mrs R. Legrand (2015), **I:** Mrs R. Legrand (2015), **REG:** Mrs R. Legrand (2015).

Fls 9–11/compact truss, tubular funnel-shaped, 65 × 42mm, with 5 wavy-edged lobes, buds pale yellow (158A), blushed deep pink (52C), opening inside white (155D), blushed deep pink (52C) around the lobe margins, most noticeably the lobes adjacent to the dorsal lobe which has well-marked speckling of strong orange-yellow (17A); stamens 80mm long; good scent. Calyx 8mm long, strong yellow-green (144B). Lvs narrowly elliptic, 105 × 40mm, attenuate at base, acute at apex, margins decurved, matt green above. Shrub 1.8 × 1.15m in 30 years. May.  
Etymology: named for the late wife of a friend of the registrant.

(r) **‘Pink Icicles’**

cv. *Elepidote rhododendron*: seedling from selfed ‘Spring Morning’.

**H:** B.N.Starling (1990), **G:** B.N.Starling (1995), **N:** B.N.Starling, **I:** B.N.Starling (1999), **REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot

tip, urceolate, 13 × 9mm, with 5 flat-edged lobes, buds cream shading to purple at tips, opening inside cream, shading to pink lobes; outside white to cream, shading to pink lobes; anthers light brown; style creamy green. Calyx 3mm long, green. Pedicels with fine hairs. Lvs elliptic, 70 × 37mm, cuneate at base, acute at apex, matt bluish green above, hairy above, with a few bristles on the veins below; petioles sparsely hairy. Upright, fairly open shrub, 1.5 × 1.0m in 10 years, with hairy stems. April to May. Hardy to at least –15C.

PC 1992.

Previously listed as *Menziesia* ‘Pink Icicles’. The epithet alludes to the superficial resemblance of the flowers to icicles with a pink tip.

(r) **‘Radiant Pink Lady’**

cv. *Elepidote rhododendron*: ‘Brookville’ × ‘Mary Garrison’ (s) × ‘Crest’ seedling (no. 2) × ‘Crest’ seedling (no. 1).

**H:** H.E.Reiley (2006), **G:** M.L.Reiley (2015), **S:** M.L.Reiley (2015), **N:** M.L. Powell (2015), **REG:** M.L.Reiley (2015).

Fls 12–14/ball truss, broadly funnel-shaped, 44 × 64mm, with 5 emarginate, wavy-edged lobes, buds light yellow-green (2C), blending to light yellowish pink (36C), opening inside brilliant greenish yellow (3A), shading to pale yellowish pink (36D) at margins, with red dorsal spots; outside light greenish yellow (3C) at base, shading to light yellowish pink (36B–C). Calyx 2mm long, strong yellow-green (145A). Lvs elliptic, 90–95 × 32–38mm, rounded at base, acute at apex, margins flat, semiglossy and moderate olive green (137A) above, hairless. Shrub 1.1 × 0.9m in 9 years. Early (early May in central Maryland, USA). Hardy to –19C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119759).

(r) **‘Reiley’s Cream Puff’**

cv. *Elepidote rhododendron*: (‘Brookville’ × ‘Mary Garrison’) × (unnamed yellow seedling (no. 9-59) × ‘Tan’) (s) × probably ‘Golden Star’ seedling × (Jalisco Group × *yakushimanum*).

**H:** H.E.Reiley (2005), **G:** M.L.Reiley (2015), **S:** S.Coble (2015), **N:** S.Coble (2015), **REG:** M.L.Reiley (2015).

Fls 12–15/ball truss, broadly funnel-shaped, 51 × 76mm, with 7 rounded, wavy-edged lobes, light greenish yellow (3D) in bud, opening inside pale yellow (158B), edged with light yellowish pink (36C), with light yellowish pink (36B) spots on the dorsal lobe; outside pale yellow (158B), edged with light yellowish pink (36C); lightly scented. Calyx 3mm long, strong yellow-green (145A). Lvs oblong, 76–90 × 44mm, rounded at base, obtuse at apex, margins flat but blade slight concave, moderate olive green (147A) and semi-glossy above, hairless. Shrub 1.4 × 2.1m in 10 years. Early (early May in central Maryland, USA). Hardy to –19C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119757).

(r) **'Rektors'**

cv. *Elepidote rhododendron*: 'Catawbiense Grandiflorum' (s) × 'Mrs P.D. Williams'.

**H:** R.Kondratovičs (1973), **G:** R.Kondratovičs (1980), **N:** R.Kondratovičs (2012), **I:** Rhododendron Nursery Babite, University of Latvia (2014),

**REG:** Rhododendron Nursery Babite, University of Latvia (2015).

Fls 14/strongly domed truss, open funnel-shaped, 55 × 70mm, with 5 wavy-edged lobes, strong reddish purple (72B) in bud, opening inside very pale purple (76C), with conspicuous deep greenish yellow (153A) spots on the dorsal lobes; outside light purple (75B); filaments light purple, anthers grey; style red-purple, with a red-purple stigma. Calyx 5mm long, light green. Lvs elliptic, 155 × 60mm, rounded at base, pointed at apex, margins flat to slightly wavy, glossy green above, hairless. Broad, bushy shrub, 1.9 × 2.1m in 20 years. Late May to early June. Hardy to -30C. Nomenclatural standard: colour print provided by the registrant (WSY0119766). Etymology: named for a rector of the University of Latvia.

(r) **'Rio Carnival'**

cv. *Elepidote rhododendron*: 'Seaview Sunset' (s) × 'Autumn Gold' × 'Orange Marmalade'.

**H:** F.Fujioka (1998), **G:** F.Fujioka (2005), **N:** F.Fujioka (2015), **REG:** F.Fujioka (2015).

Fls 20/domed truss, funnel-campanulate, 64 × 76mm, with 5 flat-edged lobes, buds moderate reddish orange (35A) at tips, shading through strong yellowish pink (35C) to moderate pink (36D) at the base; opening inside light yellowish pink (19B), shading to a 13mm marginal band of strong yellowish pink (37A), with insignificant spotting; outside strong yellowish pink (37A) at margins, shading to light orange-yellow (22B). Calyx 38mm long, strong yellowish pink (37B). Lvs elliptic, 127 × 51mm, rounded at base, obtuse at apex, margins downcurved, semi-glossy and dark bluish green (133A) above, hairless. Shrub 1.0 × 1.3m in 12 years. Late (late May in Puget Sound, Washington, USA). Hardy to -12C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119707).

(r) **'Roseum Mustila'**

Syn. of 'Kristian's Pink'.

(r) **'Roxy'**

cv. *Elepidote rhododendron*: 'Bruns Diana' (s) × 'Marrakesch'.

**H:** H.Hesse (1996), **G:** H.Hesse (2001), **N:** H.Hesse (2008), **REG:** H.Hesse (2015).

Fls 8–12/rounded, but slightly loose and flattened truss, open funnel-shaped, 42–55 × 62–80mm, with 5 wavy-edged lobes, inside opening deep pink (48A, 50B), with a hint of orange on the dorsal lobe, ageing to strong pink (50C), with cream-coloured sections; outside strong red (53C) as it starts to open, opening to deep pink (50B) to strong red (51A), ageing to deep pink (51C) along the midveins and lobe margins, light yellowish pink (27A, 36B) either side of midveins; with vivid red (45C) spotting at the base of the dorsal lobe and moderate red (47A) basal

markings; stamens pinkish white; style salmon orange at base, salmon red at apex. Calyx 2–15mm long, petaloid, strong red to deep pink and strong pink (47B, 47D and 50C). Lvs elliptic to broadly lanceolate, 80–120 × 37–46mm, rounded at base, broadly acute at apex, margins flat, glossy green above, hairless. Rounded, upright shrub, 0.9 × 0.8m in 10 years. Mid May to mid June. Hardy to -24C. Nomenclatural standard: colour print provided by the registrant (WSY0119733).

(a) **'Ryleigh Paige'**

cv. Evergreen azalea: 'Elsie Lee' (s) × 'Satellite' (Klupenger).

**H:** J.F.Klimavicz (1991), **G:** J.F.Klimavicz (1993), **N:** J.F.Klimavicz (2015), **REG:** J.F.Klimavicz (2015). Fls fully double, 2–3/ single bud inflorescence, 76mm wide, with 30 broadly rounded, overlapping and very wavy-edged lobes (the outer saucer-shaped when fully open), strong purplish pink (67D to 68B) in bud, opening inside strong purplish pink (68B), shading to deep purplish pink (68A) in centre, with a faint blotch of strong purplish red (67A); outside strong purplish pink (67D to 68B); no stamens and most fls with no pistil either. Calyx 7mm long, moderate yellow-green (138B). Lvs elliptic, 38 × 16mm, acute at apex, cuneate at base, margins flat, upper surface glossy, somewhat bullate and moderate olive green. Shrub 1.2 × 1.2m in 10 years. Midseason (mid May in Washington, D.C., USA). Hardy to -23C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a niece of the hybridiser's son-in-law.

(r) **'Spring Flare'**

cv. *Elepidote rhododendron*: 'Scintillation' (s) × 'Firestorm'.

**H:** J.Doppel (2000), **G:** J.Doppel (2004), **N:** J.Doppel (2014), **REG:** J.Doppel (2015). Fls 11/domed truss, open funnel-shaped, 38mm long, with 7 emarginate, wavy-edged lobes, buds deep red (53A), opening inside strong purplish red (58B), with a pale purplish pink (56A) flare on the dorsal lobe extending almost to the base where there is a small star of strong purplish red (71B); outside shading from strong purplish pink (55B) at base to vivid purplish red (N57A) at margins. Calyx insignificant. Lvs elliptic, 100 × 50mm, rounded at base, broadly acute at apex, margins downcurved, moderate yellow-green (137C) above, hairless. Dense shrub, 0.9 × 2.1m in 15 years. Early (May in eastern Pennsylvania, USA). Hardy to -23C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119784).

(r) **'Spring Morning'**

cv. *Elepidote rhododendron*: *benhallii* 'Buchanan's Dwarf' × *menziesii*, glaucous form.

**H:** B.N.Starling (1983), **G:** B.N.Starling (1989), **N:** B.N.Starling, **I:** B.N.Starling (1991),

**REG:** B.N.Starling (2015).

Fls pendulous, 4–6 from each of 1–5 buds at a shoot tip, urceolate, 10 × 8mm, with 5 flat-edged lobes, smoky amber in bud, opening inside cream, shading



to salmon pink on the lobes, outside amber, shading to salmon pink on the lobes; anthers dark brown; style cream. Calyx 2mm long, green. Pedicels with fine hairs. Lvs elliptic, 43 × 23mm, cuneate at base, acute at apex, matt and glaucous above, hairy above, with a few bristles on the veins below; petioles sparsely hairy. Compact, slightly spreading habit, 1.5 × 1.0m in 10 years, with hairy stems. May to June. Hardy to at least -15C.

AM 1992.

Previously listed as *Menziesia* 'Spring Morning'. The epithet alludes to the effect of the flower colour seen against the glaucous foliage.

(r) 'Standing Ovation'

cv. Elepidote rhododendron: 'Orange Prelude' (s) × Fortune Group × ('Berg's Yellow' × *proteoides*).

H: J.Barlup (2003), G: J.Barlup (2009), N: J.Barlup (2015), REG: J.Barlup (2015).

Fls 16/ball truss, funnel-campanulate, 51 × 76mm, with 7 wavy-edged lobes, buds strong red (53B), opening inside pale yellow (11C), blending to strong pink (52D) at margins, with a small, deep red (53A) basal circle, outside pale yellow (11C), blending to strong pink (52D) at margins, with a strong pink (52D) central rib line on each lobe; not scented. Calyx 16mm long, pale yellow (11C), with two 6mm long dorsal flares. Lvs elliptic, 152 × 73mm, broadly acute at apex, rounded at base, margins flat, moderate olive green (147A) above, hairless; new growth deep bronze. Open shrub, 0.9 × 0.9m in 6 years. Midseason (late April in Seattle, Washington, USA). Hardy to -15C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119845).

(r) 'Steve's Sunrise'

cv. Elepidote rhododendron: 'Janet Blair' × ('Evening Glow' × 'Crest seedling (no. 2)') (s) × unnamed yellow seedling × ('Evening Glow' × 'Crest' seedling (no. 2)).

H: H.E.Reiley (2004), G: M.E.Reiley (2015), S: M.E.Reiley (2015), N: S.E.Reiley (2015), REG: M.E.Reiley (2015).

Fls 8-11/ball truss, broadly funnel-shaped, 51 × 76mm, with 5-7 broadly acute, wavy-edged lobes, buds moderate purplish red (186A) to dark purplish pink (186C), opening inside pale yellowish pink (36D), with light orange-yellow (16C) at base and moderate purplish red (186B) at margins, and moderate reddish orange (35A) speckling on the dorsal lobe; outside dark purplish pink (186C), with moderate purplish red (186A) midveins. Calyx 3mm long, strong yellow-green (145A). Lvs elliptic, 64-76 × 32-38mm, rounded at base, broadly acute at apex, margins downcurved, matt and moderate olive green (137A) above, hairless. Shrub 1.1 × 1.1m in 11 years. Early (early May in central Maryland, USA). Hardy to -19C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119755). Etymology: named for the son (S.E.Reiley) of the hybridiser and registrant.

(r) 'Storm'

cv. Elepidote rhododendron: 'Spring Morning' × *multiflorum* var. *purpureum*.

H: B.N.Starling (2002), G: B.N.Starling (2005), N: B.N.Starling (2010), I: B.N.Starling (2012), REG: B.N.Starling (2015).

Fls pendulous, 4-6 from each of 1-5 buds at a shoot tip, urceolate, 15 × 9mm, with 5 flat-edged lobes, muddy rose in bud, opening inside cream tinged with pink and with red lobes, outside rose shaded with grey, with red lobes; anthers brown; style creamy white; not scented. Calyx 4mm long, dark purple-black. Pedicels with fine hairs. Lvs broadly ovate, 40 × 27mm, cuneate at base, rounded at apex, margins flat, matt green above, sparsely hairy. Shrub with a fairly open, branching habit, 1.0 × 1.0m in 10 years. May to June. Hardy to at least -15C.

Previously listed as *Menziesia* 'Storm'. Etymology: the epithet relates to the colour of the flowers, which is reminiscent of storm clouds.

(v) 'Sushmitam Rouse'

cv. Lepidote rhododendron: *sessilifolium* (s) × 'Clare Rouse'.

H: A.Rouse (1998), G: A.Rouse (2006), N: A.Rouse (2015), REG: A.Rouse (2015).

Fls 6-9/open truss, tubular funnel-shaped, 50-53 × 50-52mm, with 5 slightly wavy-edged lobes, light green in bud, opening both inside and out light greenish yellow (4B), becoming brilliant yellow as the flower matures, unmarked; style and filaments coloured as the corolla; not scented. Calyx and pedicels a uniform light yellow-green (154D). Calyx less than 1mm long. Lvs elliptic, 50-100 × 15-35mm, tapering at base, acute to acuminate at apex, margins slightly wavy, concave, glossy green, with tiny, sparse scales (not clearly visible to the naked eye). Shrub with an open habit, 1.0 × 0.8m in 17 years. May (i.e. autumn in Australia).

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119782).

Etymology: named for the registrant's sister.

(r) 'Tamera'

cv. Elepidote rhododendron: *yakushimanum* 'North Island' (s) × *grande*.

H: P.Smith (2001), G: P.Smith (2012), N: P.Smith (2013), REG: P.Smith (2015).

Fls 22-28/domed truss, funnel-campanulate, 51 × 51mm, with 7 wavy-edged lobes, pinkish white in bud, opening inside white, with moderate pink (49B) veins and a strong yellowish pink (37B) flare extending 13mm from the base on the dorsal side of the corolla; outside white. Calyx 2mm long, green. Lvs elliptic, 152 × 57mm, cuneate at base, broadly acute at apex, margins downcurved, glossy and dark green (136A) above, tomentose above when young, the hairs persistent below and white at first, ageing to beige. Dense shrub, 1.4 × 0.9m in 10 years. Early (March in northern California, USA). Hardy to -8C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119709). Etymology: named for a daughter of the registrant.

(a) 'Tarleena'

cv. Deciduous azalea: 'Directeur Charles Baumann' (s) × Choptank River Hybrid × 'Maori'.

**H:** A.Väinölä (1989), **G:** University of Helsinki and Arboretum Mustila, **N:** A.Väinölä (2010), **I:** MTT Agrifood Research Finland (2012), **REG:** University of Helsinki (2015).

Fls 7–10/truss, narrowly funnel-shaped, 43–48 × 38–42mm, with 5 flat-edged lobes, buds vivid red (45B, 46B), opening inside deep to strong pink (45A,B,C, 49A, 50C, 51C), the colour varying from flower to flower with age, outside deep pink (51B, 52B); middle of the dorsal lobe vivid orange-yellow (21A, 23A).

Calyx 2mm long, light green. Lvs mostly elliptic, 90 × 23–25mm, cuneate at base, acute at apex, margins flat, matt green above, hairy. Shrub 1.65–1.75 × 1.7–2.0m in 15 years. Late (mid June in Finland). Hardy. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY019778).

**Etyymology:** named for a gnome in a Finnish fairy tale by E. Karjalainen (2003). First published in *Sorbifolia* 43(1):34–38 (2012).

(r) ‘**Teshio**’

cv. *Lepidote rhododendron*: Selection from *hypoleucum* (syn. *Ledum palustre* var. *diversipilosum*).

**G:** B.N.Starling (1991), **N:** B.N.Starling,

**I:** B.N.Starling (1994), **REG:** B.N.Starling (2015).

Fls c.50/composite truss (which is up to 10cm across and formed from the flowers of 3–5 buds), star-shaped, with 5 petals, 20mm wide, white in bud and throughout on opening; anthers and style white. Calyx 2mm long, green. Lvs lanceolate, 50 × 10mm (sometimes up to 100mm long), cuneate at base, margins recurved, dark green above, glaucous below and with fawn hairs when young, which age to dark brown. Shrub 1.0 × 1.0m in 10 years. Hardy to at least –15C.

PC 1992.

Formerly listed as a cultivar of *Ledum palustre* var. *diversipilosum*. **Etyymology:** derived from seed collected by the registrant, found on the territory covered by the Teshio Forestry Research Station in northern Hokkaido, Japan during an AGS expedition in 1988.

(r) ‘**Teterevi Latvijai**’

cv. *Elepidote rhododendron*: ‘*Britannia*’ (s) × ‘*Mrs R.G. Shaw*’.

**H:** R.Kondratovičs (1974), **G:** R.Kondratovičs (1980), **N:** R.Kondratovičs (2015), **REG:** Rhododendron Nursery Babīte, University of Latvia (2015).

Fls 17/strongly domed truss, funnel-shaped, 55 × 65mm, with 5 wavy-edged lobes, moderate purplish red (58A) in bud, opening inside light purplish pink (73C), with dark red (59A) spots on the dorsal lobes, outside deep purplish pink (73A); filaments red-purple, anthers beige; style red-purple, with a red stigma. Calyx 2mm long, light purple. Lvs elliptic, 130 × 60mm, rounded at base, pointed at apex, margins flat, matt green above, hairless. Bushy shrub, 2.4 × 1.9m in 25 years. Late May to early June. Hardy to –25C.

Nomenclatural standard: colour print provided by the registrant (WSY0119767). **Etyymology:** named for the creators of the Teterev family’s Charity Foundation.

(r) ‘**Watermelon Sky**’

cv. *Elepidote rhododendron*: ‘*Sumner’s Orange*’ (s) × ‘*Double Eagle*’ (Korth).

**H:** P.Smith (2001), **G:** P.Smith (2012), **N:** P.Smith (2013), **REG:** P.Smith (2015).

Fls 6–8/domed truss, funnel-campanulate, 51 × 89mm, with 5–6 flat-edged lobes, strong red (46A) in bud, opening inside deep pink (51B), with strong red (51A) margins and a 13mm strong red (53B) spot at the base, surmounted by a vivid orange (28B) blotch on all lobes; outside deep pink (51B). Calyx up to 13mm long, deep pink (50B). Lvs elliptic, 127 × 64mm, rounded at base, broadly acute at apex, margins flat, dull and moderate olive green (146A) above, hairless. Shrub 0.9 × 0.9m in 10 years. Early (February to March in northern California, USA). Hardy to –8C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119711).

(r) ‘**Whisper White**’

cv. *Elepidote rhododendron*: ‘*Nancy Evans*’ × *bureavii* (s) × *yakushimanum* ‘*Koichiro Wada*’ × *pachysanthum*.

**H:** F.Fujioka (2003), **G:** F.Fujioka (2008),

**N:** F.Fujioka (2014), **REG:** F.Fujioka (2015).

Fls 18/ball truss, funnel-campanulate, 51 × 64mm, with 5 wavy-edged lobes, buds light pink (49C), opening inside white (155D), with a vivid red (44A) flare and spots, which extend 19mm on the dorsal side, but less on the others; outside white (155D), with 13mm of pale pink (49D) at the base. Calyx insignificant. Lvs elliptic, 140 × 64mm, rounded at base, broadly acute at apex, margins flat, dull and dark yellowish green (139A) above, hairy on both sides (less so above), the hairs creamy white (159D) above, greenish white (157D) below at first and ageing to moderate orange-yellow (165C) below. Shrub 0.9 × 0.9m in 10 years. Midseason (late April in Puget Sound, Washington, USA). Best grown in slight shade. Hardy to –12C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119713).

(r) ‘**Will Corey**’

cv. *Elepidote rhododendron*: unnamed seedling (Morris 452) (s) × *fortunei*.

**H:** P.B.Moser (2007), **G:** P.B.Moser (2014),

**N:** P.B.Moser (2014), **REG:** P.B.Moser (2015).

Fls 12–13/ball truss, open funnel-shaped, 51 × 90mm, with 5–7 rounded, flat-edged lobes, strong purplish red (63B) in bud, opening inside strong purplish pink (68B), with small, twin, basal flares of strong greenish yellow (151B), outside deep purplish pink (68A); slightly scented. Calyx 2mm long, strong yellow-green (144C). Lvs elliptic, 127 × 43mm, rounded at base, broadly acute at apex, margins flat but lamina slightly concave, moderate olive green (146A) above, hairless. Shrub 0.8 × 0.9m in 8 years. Midseason (May in Philadelphia, Pennsylvania, USA). Hardy to –18C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119783).

**Etyymology:** named for the deceased son of a friend of the registrant.

# Corrections to the *International Rhododendron Register and Checklist (2004)*

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to its subsequent Supplements. The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made. Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

**'Crème Brûlée'** (Second Supplement, p.10)

This should be treated as a synonym of 'Lemon Meringue' as it is now known to be a later naming of the same plant.

**'Cream Delight'** (p.293)

The roles entries should be amended to B.W. Campbell throughout, not Sir Ilay Campbell.

**'Dame Kiri Tekanawa'** (p.310)

This should be given as 'Dame Kiri Te Kanawa' (*ie* four words).

**'Frank Mossman'** (p.469)

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119691).

**'Heritage Snow Ruby'** (Seventh Supplement, p.13)

This should be treated as a synonym of 'The Morning After' as it is now known to be a later naming of the same plant.

**'King Elliott'** (p.722)

The roles entries should be amended to E.W.E. Butcher, not E.W.E. Butler.

**'Tatum's Deep Pink'** (Fifth Supplement, p.29)

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119710).

**'Tatum's Yellow Standard'** (Third Supplement, p.23)

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0119712).

# Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

Arboretum Mustila Elimäki, Finland	Hahkala, R. Finland
Barlup, J. Bellevue, Washington, USA	Hesse, H. Hamersen, Germany
Baumschule H. Hachmann, Barmstedt, Germany	Hübbers, K.-H. & Mrs L. Kranenburg, Germany
Beaudry, N. Bethesda, Maryland, USA	Humphries, K.F. USA
Begg, L. Australia	Hyde, W.G.T. Ferndown, Dorset, England
Brindabella Country Gardens Toowoomba, Queensland, Australia	Johnson, P. Fort Bragg, California, USA
Charvet, L. Fort Bragg, California, USA	Klimavicz, J.F. Vienna, Virginia, USA
Coble, S. Gettysburg, Pennsylvania, USA	Kondratovičs, R.K. Rīga, Latvia
Cox, P.A. & K.N.E. Glendoick Gardens, Glencarse, Perthshire, Scotland	Lansdowne, Marquis of Bowood House, Calne, Wiltshire, England
Delano, J. Duxberg, Massachusetts, USA	Legrand, Mrs R. West Moors, Dorset, England
Doppel, J. Lenhartsville, Pennsylvania, USA	Millais, E.G. & D. Millais Nurseries, Farnham, Surrey, England
Forest Lane Nursery Fort Bragg, California, USA	Moser, P.B. Bryn Mawr, Pennsylvania, USA
Fujioka, F. Freeland, Washington, USA	MTT Agrifood Research Finland Vihtavuori, Finland
Furman, R.A. Brewster, Massachusetts, USA	Powell, M.L. Woodsboro, Maryland, USA
Gehnrich H.C. Huntingdon, New York, USA	Reiley, H.E. & Mrs M.L. Woodsboro, Maryland, USA
Glendoick Gardens Glencarse, Perthshire, Scotland	Rhododendron Nursery Babite, University of Latvia Rīga, Latvia
Goossens, J. Lochristi, Belgium	Rouse, A. Hawthorn East, Victoria, Australia
Gray, J.C. Toowoomba, Queensland, Australia	Singing Tree Gardens McKinlayville, California, USA
Gustavson, J. Victoria, British Columbia, Canada	Smith, P. Santa Monica, California, USA

Starling, B.N.  
Drewsteignton, Exeter, Devon, England

Stewart, R.  
Springfield, Virginia, USA

Stokes, I.T.  
Landeilo, Carmarthenshire, Wales

Theqvist, K.  
Turku, Finland

Trujillo, R. & S.  
Fort Bragg, California, USA

University of Helsinki  
Finland

Väinölä, A.  
Vantaa, Finland

Van Veen Nursery  
Portland, Oregon, USA

Wallace, D.S.  
Singing Tree Gardens, McKinlayville,  
California, USA

Westhoff, J.  
Bremen, Germany

White, G.P.  
Kingham, Oxfordshire, England

Wüstemeyer, W.  
Schermbeck, Germany





ISBN 9781907057724



9 781907 057724