



Inspiring everyone to grow

RHS

# The International Rhododendron

## Register & Checklist (2004)

*Fourteenth Supplement*



© 2019 The Royal Horticultural Society  
80 Vincent Square, London SW1P 2PE, United Kingdom  
[www.rhs.org.uk](http://www.rhs.org.uk)

International Rhododendron Registrar:  
Sharon McDonald  
Email: [sharonmcdonald@rhs.org.uk](mailto:sharonmcdonald@rhs.org.uk)

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057953

Printed and bound in the UK by Page Bros, Norwich (MRU)

The previous supplement (Thirteenth Supplement) was published on 29 May 2018.

*Cover: Rhododendron 'Fireball', by Elsie Katherine Kohnlein*  
Image courtesy of RHS Lindley Library

# THE INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST (2004)

## *Fourteenth Supplement*

Names registered, January–December 2018	page 2
Corrections	page 22
Names and addresses	page 23

## Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first thirteen Supplements included names registered from 1 January 2003 to 31 December 2017. This *Fourteenth Supplement* includes all those registered in 2018.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction.

It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea
(az)	azaleodendron
cv.	cultivar
G	grown to first flower by
gp	Group
H	hybridised by
I	introduced by
N	named by
(r)	rhododendron
R	raised by
REG	registered by
S	selected by
(s)	seed parent
(v)	vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995, 2001 & 2015) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website (<https://www.rhs.org.uk/plants/pdfs/plant-registration-forms/rhododendron-name-registration.PDF>) or from one of the national or regional registrars as follows:

<b>North America</b>	Mr Michael Martin Mills, 632 Burnham Road, Philadelphia, Pennsylvania PA19119-3510, USA (Email: <a href="mailto:arsregistrar@gmail.com">arsregistrar@gmail.com</a> )
<b>Australia</b>	Mrs Lesley Eaton, PO Box 597, Snug, Tasmania 7054, Australia (Email: <a href="mailto:lesley.eaton@bigpond.com">lesley.eaton@bigpond.com</a> )
<b>New Zealand</b>	Mr Brian J. Coker, Stonecrop, 8B Barnsley Crescent, West Melton, New Zealand 7618 (Email: <a href="mailto:b.hcoker@xtra.co.nz">b.hcoker@xtra.co.nz</a> )
<b>Japan</b>	Dr Katsuhisa Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

**Dr A.C. Leslie, International Rhododendron Registrar, retired in June 2018.**

**New registrations should be sent to:** Sharon McDonald, RHS Garden Wisley, Woking, Surrey, GU23 6QB, UK  
(Email: [sharonmcdonald@rhs.org.uk](mailto:sharonmcdonald@rhs.org.uk))

## Names registered January–December 2018

(r) '24 Caret'  
Syn. of '24 Karat'

(r) '24 Karat'

cv. Lepidote rhododendron: *keiskei* (s) × Nearing's form of *cinnabarinum* subsp. *xanthocodon*

**H:** C.R. & V. Haag, **G:** C.R. & V. Haag, **N:** C.R. & V. Haag, **REG:** M.M. Mills (2018)

Fls 8–10/flat truss (3–5/bud, 1–3 buds/truss), funnel-shaped, 32 × 38mm, with 5 acute, reflexed, pinched and minutely very wavy-edged lobes, brilliant yellow-green (149B) in bud, opening pale greenish yellow (1D), with deep greenish yellow (153A) spots and throat below the dorsal lobe; outside pale greenish yellow (1D); filaments pale greenish yellow (2D), anthers light yellow (12C); style pale greenish yellow (2D), with a strong yellowish pink (32C) stigma; style goes pink after pollination. Calyx insignificant. Lvs elliptic, 47 × 16mm, rounded at base, acute at apex, margins flat, midrib depressed, moderate yellow-green (137C) above, with scales below and with minute hairs below and on petioles. Shrub 0.3–0.4 × 0.6m in 8–10 years. Early (early April in Philadelphia, Pennsylvania, USA). Hardy to at least –17C.

Syn: '24 Caret', 'Twenty Four Carat'

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125616). Originally listed by the Haags as '24 Caret' but this is treated here as an unintentional error (in a list with other spelling errors), as it is clear that the reference was to gold and overall excellence, rather than the printing term caret. The American spelling of "karat" is deliberate. Date of origin is uncertain but perhaps in the 1970s or 1980s.

(r) 'Absolute Citron'

cv. Elepidote rhododendron: 'Ada Lohr' (s) × 'Goldkrone'

**H:** R.R. Rabideau (1996), **G:** R.R. Rabideau (1999), **N:** R.R. Rabideau (2006), **I:** RareFind Nursery (2008), **REG:** R.R. Rabideau (2018)

Fls 13/domed truss, funnel-shaped, 51 × 76mm, with 5 rounded, wavy-edged lobes (some emarginate, others pinched apically to give an emarginate appearance), brilliant greenish yellow (3B) in bud, opening inside light greenish yellow (3C), with brilliant greenish yellow (3B) in centre of dorsal lobes and on the midveins, and with a dorsal blotch of strong red (46A) speckles; outside light greenish yellow (3C), with the speckles showing though; style and filaments yellowish; ovary with dense, stalked, red glands. Calyx 3mm long, light yellow-green (145C), with scattered, stalked, red glands. Lvs elliptic, 102 × 44mm, rounded at base, obtuse at apex, margins flat, matt and deep yellowish green (darker than 141A) above. Shrub 0.9 × 1.5m in 12 years. Mid-season (mid May in central New Jersey, USA). Hardy to –29C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125617). First published in RareFind Nursery cat. 2008.

(a) 'Alan Leslie'

cv. Deciduous azalea: *prunifolium* (s) × *viscosum* f. *rhodanthum*

**H:** E.G. Millais (1999), **G:** E.G. Millais (2002), **N:** D. Millais (2018), **I:** Millais Nurseries (2018), **REG:** D. Millais (2018)

Fls 9–10/rounded truss, very narrowly tubular funnel-shaped (opening flat-faced), 40–50 × 35–40mm, with 5 flat lobes, buds deep purplish pink (54B), opening inside deep purplish pink (54B) at margins, paling to strong pink (54C) at centre; outside tube and midrib deep purplish pink (54B), shading to strong pink (54C); filaments bright pink, paling to base, anthers brown; style bright pink, with a creamy white stigma; scented. Calyx 3–5mm long, very pale pink/white. Lvs elliptic, 65–80 × #

19–25mm, shortly attenuate at base, broadly acute at apex, with margins decurved, slightly glossy above, hairs on midrib only, becoming slightly twisted with maturity. Shrub 0.75 × 0.75m in 10 years. Late July. Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125566).

Etymology: named for the recently retired International Rhododendron Registrar.

(r) 'Anta'

cv. Elepidote rhododendron: 'Edmond Amateis' (s) × *macabeanum*

**H:** T. Knights (1992), **G:** J. Jäck (2010), **N:** J. Jäck (2015), **REG:** J. Jäck (2018)

Fls single, 11–16/roundish truss, funnel-campanulate, 55 × 55–65mm, with 7 wavy-edged lobes, buds light pink, opening inside white, with red basal markings and some spots flared out from the base of the throat; outside white. Lvs oblong, 140–180 × 60–80mm, broadly acute at apex, matt green above, with pale whitish hairs when young, aging to greyish beige (but only as a light indumentum). Shrub 2.2 × 2.0m in 25 years. Late April to early May (in northern Germany). Hardy to –20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125504).

Etymology: Anta is the Portuguese word for tapir.

(r) 'Arctic Morn'

cv. Elepidote rhododendron: 'Sun Blush' (s) × 'Golden Mist'

**H:** J. Barlup (2009), **G:** J. Barlup (2015), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 19/ball truss, broadly funnel-shaped, 51 × 76mm, with 5 rounded, wavy-edged lobes, pale yellow (11C) in bud, opening inside pale yellow-green (4D), with light greenish yellow (4C) in the dorsal section of the throat, and with a prominent deep greenish yellow (153A) spotted blotch on the dorsal lobe and on flanking halves of the adjacent lobes; outside pale yellow-green (4D), with light greenish yellow (4C) on the dorsal lobes from base to 6mm from margins; filaments yellowish, anthers light brownish; pistil yellowish. Calyx 25mm long, pale yellow-green (4D), shading to light greenish yellow (4C). Lvs elliptic, 127 × 44mm, rounded at base, broadly acute at apex, margins flat,

semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 8 years. Mid-season (mid May in Seattle area, Washington, USA). Hardy to –18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108639).

(r) **'Ariadne Grace'**

cv. *Elepidote rhododendron*: 'Apricot Fantasy' × 'Phipps Yellow' (s) × 'Virginia Delp'  
**H:** G. Woodard (2002), **G:** G. Woodard (2009),  
**N:** M. Calvo-Platero (2011), **REG:** Howard Phipps Jr Estate (2018)  
Fls 13/domed truss, funnel-shaped, 51 × 76mm, with 5 rounded, very wavy-edged lobes, white in bud, opening inside white, with vivid greenish yellow (2A) dorsal area from base into the lobes, surmounted by a conspicuous, strong orange (170A) spotted blotch and flanking areas of spots; outside white; filaments whitish; style whitish, with a pale yellow stigma. Lvs elliptic, 127 × 38mm, cuneate at base, broadly acute at apex, margins flat, matt and moderate olive green (137A) above. Shrub 3.0 × 3.0m in 10 years. Mid-season. Hardy to –21C.

Nomenclatural standard: colour print provided by the registrant (WSY0108641). Etymology: named for the wife of the nominant, the naming rights obtained in an auction at Old Westbury Gardens, New York, USA.

(a) **'Artistic Meg'**

cv. Evergreen azalea: parentage unknown  
**H:** W.G.T. Hyde (1960s), **G:** W.G.T. Hyde (1970s),  
**N:** Mrs R.G. Legrand (2017), **I:** Mrs R.G. Legrand (2018), **REG:** Mrs R.G. Legrand (2018)  
Fls semi-double, with a cluster of short petaloids in the centre, fluted open funnel-shaped, 57 × 30mm, with 5 wavy-edged main lobes, strong purplish red (60D) in bud, opening strong purplish red (64C), with irregular and sporadic whitish marbling (between 62D and 155D); outside vivid purplish red (57C); with slight spotting of deep red (60A) on the three dorsal lobes. Calyx irregular, 5–9mm long, moderate yellow-green (146B). Lvs elliptic, 39 × 18mm, cuneate at base, acute at apex, margins flat, matt green above, hairless. Shrub 1.15 × 1.07m in c. 40 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY0108688). Etymology: named for the botanical artist Meg Mulcahy.

(r) **'Blueberry Kiss'**

cv. *Elepidote rhododendron*: 'Midnight Mystique' (s) × 'Stadt Delmenhorst'  
**H:** Holger Hachmann (2005), **G:** Baumschule H. Hachmann, **N:** Holger Hachmann, **I:** Baumschule H. Hachmann, **REG:** O. Tschetsch (2018)  
Fls 17–20/slightly flattened ball truss, rotate, 35 × 15mm, with 5 wavy-edged, sometimes emarginate, lobes, buds deep reddish purple (77A), opening inside very pale purple (76D), with edge and 5mm margin of strong purple (80B) and with dark purple (79A) spots on the dorsal and adjacent lobes; outside light purple (78C); nectaries moderate

purplish red (58A); 10 stamens, filaments very pale purple (76D), anthers pale greenish yellow (10D); style greenish white (157D), with light yellow-green (154D) at base, stigma vivid yellow-green (154A); not scented. Calyx 2mm long. Pedicels 30mm long, moderate purplish red (58A) and strong yellow-green (144B). Lvs elliptic, 110 × 45mm, rounded at base, broadly acute at apex, with wavy and upcurved margins, matt, moderate olive green (146A) above, moderate yellow-green (146C) below, hairless. Shrub 1.3 × 1.6m in 10 years. Early to mid May (in northern Germany). Hardy to –22C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125581).

(r) **'Camdeboo'**

cv. *Elepidote rhododendron*: Albatross Group (s) × 'Britannia'  
**H:** F. Schnupper (2002), **G:** F. Schnupper,  
**N:** F. Schnupper, **I:** Schnupper Baumschule (2018),  
**REG:** O. Tschetsch (2018)

Fls single, 12/domed truss, funnel-shaped, 60 × 70mm, with 7 wavy-edged lobes, buds strong purplish red (58B–C), opening inside light purplish pink (62C), lightly marked with strong purplish pink (62A), the dorsal and adjacent lobes heavily marked with strong purplish red (58B) spots, with a moderate yellow (163C) blotch in throat; outside light purplish pink (62A), lightly marked with strong purplish red (62A); 13–14 stamens, filaments light yellow-green (150D), anthers moderate orange (167A); style brilliant yellow-green (150C), with strong yellow-green (144B) at base, stigma brilliant yellow-green (150B); ovary strong yellow-green (144A). Calyx 1mm long, dark red (187B). Pedicels 25mm long, strong yellow-green (144B), with touches of dark red and deep red (185A & 187A). Lvs elliptic, 125 × 40mm, rounded at base, broadly acute at apex, margins wavy and decurved, matt, moderate olive green (137A) above, light yellow-green (144D), with brilliant yellow-green (151D) midrib below, hairless; petioles 35mm long, brilliant yellow-green (151D). Shrub 1.0 × 1.2m in 10 years. Early to late May (in northern Germany). Hardy to –22C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125545). Etymology: named after Camdeboo National Park, South Africa.

(r) **'Caroline Darvall'**

cv. *Elepidote rhododendron*: parentage unknown  
**H:** N. de Rothschild (1994), **G:** Exbury Gardens,  
**N:** N. de Rothschild (2017), **REG:** T. Clarke (2018)  
Fls 12/lax truss, tubular funnel-shaped, 60–70 × 70–120mm, with 5–6 wavy-edged lobes, strong yellowish pink (43D) in bud, opening inside pale purplish pink (56B), fading to pale purplish pink (56D); outside moderate purplish pink (65A), fading to pale purplish pink (65C); throat moderate orange-yellow (168D), most extensively on the dorsal side, with brown speckles; style lime green; ovary glandular; scented. Calyx prominent, irregularly lobed, 20–30mm long, pale purplish pink (65C), fading to moderate orange-yellow (168D) and with

brown speckles. Pedicels glandular hairy.  
Lvs oblanceolate, 110 × 35mm, cuneate at base, acute at apex, margins wavy-edged, hairless. Shrub 2.0 × 1.5m in 20 years. Mid April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125618).  
Etymology: named for the wife of the nominant.

(r) 'Chloë de Rothschild'

cv. Elepidote rhododendron: *faberi* (s) × 'Sunset over Harkwood'

**H:** N. de Rothschild (1994), **G:** Exbury Gardens (2017), **N:** N. de Rothschild (2017), **REG:** T. Clarke (2018)

Fls 8/large, rounded truss, funnel-shaped, 55 × 95mm, with 6 wavy-edged lobes, buds strong pink (50C), opening inside pale yellow (161D), with pale pink (49D) outer edges; outside moderate pink (51D); with pale yellow-green veining towards centre of flower and pale pink veining towards the margins (all darker on the outside); pale lime green style and stigma. Calyx 3–9mm long, pale yellow (161C). Lvs elliptic, 145–165 × 58–68mm, oblique at base, obtuse at apex, margins wavy and upcurved, matt green above, hairless. Upright shrub. May. Hardy in southern England.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125624).

Etymology: named for the daughter of the nominant.

(r) 'Chris's Cream'

cv. Elepidote rhododendron: 'Sunspray' (s) × 'Sunspray'

**H:** C. Mugridge (1999), **G:** C. Mugridge (2015), **N:** Christopher Mugridge (2018), **I:** Millais Nurseries (2018), **REG:** C. Mugridge (2018)

Fls 18–20/ball truss, broadly funnel-shaped, 70 × 140mm, with 7 wavy-edged lobes, buds pale creamy yellow, with touches of pink towards base, opening inside and outside pale yellow-green (4D); anthers brown; pistil green; slightly scented. Calyx 5–15mm long, pale green, pointed. Lvs elliptic, 150 × 60mm, cordate at base, broadly acute at apex, margins flat, with a satin-like upper surface, hairless. Shrub 1.0 × 1.0m in 19 years. Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125557).

Etymology: named by and for the son of the registrant.

(r) 'Christine Dohnau'

cv. Elepidote rhododendron: 'Banderilla' (s) × *yakushimanum* 'Koichiro Wada'

**H:** B. Wickhorst (2007), **G:** Baumschule H. Hachmann, **N:** C. Dohnau (2015), **I:** Baumschule H. Hachmann (2019), **REG:** O. Tschetsch (2018)

Fls 9–12/domed truss, campanulate, with 5 very wavy-edged, emarginate lobes, buds moderate red (47A), with light orange (29B) at base, opening inside brilliant yellow (9C), with deep pink (51B) margins, 8–9mm wide and with inconspicuous vivid yellow (15A) markings on the dorsal lobe; outside brilliant yellow (9C), with deep pink (51B) margins, 8–9mm wide; 10 stamens, filaments pale yellow-green (4D),

anthers moderate orange (167A); style moderate yellow (160A), stigma brilliant greenish yellow (1A); ovary strong yellow-green (143A), with dense, short white hairs; not scented. Calyx 1mm long, strong yellow-green (144A). Pedicels 37mm long, strong greenish yellow (151A/153B). Lvs elliptic, 75 × 30mm, cuneate at base, broadly acute at apex, with wavy and decurved margins, moderate olive green (147A) above and below, with strong yellow-green (145A) along the midrib on the underside; silvery hairs when young, becoming moderate brown (165A); petioles 15mm long, strong yellow-green (144A). Shrub 0.65 × 0.95m in 10 years. Early to mid May (in northern Germany). Hardy to –22C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125585).

Etymology: named by and for the Norddeutscher Rundfunk presenter.

(a) 'Coble's Edna Marie'

cv. Evergreen azalea: 'Karens' (s) × 'Girard's Fuchsia'

**H:** S. Coble (2010), **G:** S. Coble (2013), **N:** S. Coble (2017), **REG:** S. Coble (2018)

Fls 4/terminal shoot, calyx petaloid, with one lobe larger than the others, open funnel-shaped, 25 × 32mm, with 5, inner, wavy-edged lobes and 5, outer, emarginate lobes, buds vivid purple (80A), opening inside vivid reddish purple (78A), with a freckled deep reddish purple (77A) blotch; outside vivid reddish purple (78A), with moderate reddish orange (41C) at the base of the midrib; stamens and style vivid reddish purple (78A). Calyx 1mm long, vivid reddish purple (78A), and pale greenish yellow (10D). Lvs elliptic, 25 × 13mm, cuneate at base, acuminate at apex, with upcurved margins, moderate olive green (137A), darker in autumn, hairless. Shrub 0.3 × 0.6m in 8 years. Mid-season (early May in central Pennsylvania). Hardy to –23C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125569).

Etymology: named for the mother (Edna) and the sisters (Edna & Marie) of the registrant.

(r) 'Colin's 75'

cv. Elepidote rhododendron: un-named seedling (s) × 'Kluis Sensation'

**H:** C. Mugridge (2001), **G:** C. Mugridge (2017), **N:** C. Mugridge (2018), **I:** Millais Nurseries, **REG:** C. Mugridge (2018)

Fls 15–16/ball truss, broadly funnel-shaped, 40 × 90mm, with 5 wavy-edged lobes, buds deep red, opening inside deep purplish pink (67C), with a dense blotch and spots of deep purplish red (59B) on the dorsal lobe; outside strong purplish pink (63C); anthers brown; pistil pink; not scented. Calyx 3–7mm long, reddish brown. Lvs elliptic, 140 × 50mm, rounded at base, acuminate at apex, margins flat, opening lime green, becoming dark green, glossy above, hairless. Shrub 1.0 × 1.0m in 17 years. Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125558).

Etymology: named to mark the Registrant's 75th birthday.

(r) **'David Shelley'**

cv. Elepidote rhododendron: *yakushimanum* (s) × 'Purple Splendour'  
**H:** E. de Rothschild, **G:** Exbury Gardens, **N:** Lionel de Rothschild (2018), **REG:** T. Clarke (2018)  
Fls 17/domed truss, funnel-shaped, 52 × 72mm, with 5 wavy-edged lobes, vivid reddish purple (78A) in bud, opening inside very pale purple (76D) on outer edges and paling towards the centre; outside very pale purple (76C), with a slightly darker strip running down the centre of the lobes; dorsal lobe with strong purplish red (72A) spots extending down into the throat; stigma reddish pink; ovary glandular. Calyx 1mm long, pale yellow-green (195D). Lvs oblanceolate, 150 × 58mm, obtuse at base, acute at apex, margins decurved, matt green above, with a slightly woolly, moderate orange-yellow (164B) indumentum below, which disappears with age or if persisting becomes moderate orange (166D). Shrub 2 × 2m in 30 years. May. Hardy to at least -7C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125628).  
Etymology: named for a long-serving estate employee of Exbury Gardens.

(a) **'DD Gold'**

cv. Deciduous azalea: selection from wild-collected seed of *occidentale* (collection data lost)  
**G:** D. Denkers, **S:** H.R. Helm (2012), **N:** H.R. Helm (2012), **REG:** H.R. Helm (2018)  
Fls single, 19/ball-shaped truss, tubular funnel-shaped, 50 × 60mm, with 5 acute, very frilled lobes, white, with pale pink stripes in bud, opening inside white, with a large vivid yellow (15A) blotch on the dorsal lobe and with vivid yellow (15A) streaks on all other lobes, occasionally with pale pink midvein or accent at lobe tip; outside white, with pink tube; filaments, anthers and style white, stigma green; strongly scented. Calyx 3mm long, green. Lvs obovate, 40–70 × 15–30mm, cuneate at base, broadly acute at apex, margins flat, matt, light green, with some hairs, white to colourless, principally below. Shrub 2.4m high in 15 years. Mid-season (late May in Seattle, Washington, USA).  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125498).  
Etymology: incorporates the initials of the first grower.

(a) **'Dene Mathews'**

cv. Deciduous azalea: *flammeum*, open-pollinated  
**S:** V. Bush (2010), **N:** Huntsville Madison County Botanical Garden Society Inc. (2018),  
**I:** E. Somerville (2015), **REG:** Huntsville Madison County Botanical Garden Society Inc. (2018)  
Fls single, 7–9/terminal shoot, tubular funnel-shaped, 53 × 51mm, with 5–6 broadly acute, wavy-edged lobes, strong purplish red (58C) in bud, opening strong purplish pink (67D), with a modest, vivid yellow (16A) blotch on the dorsal lobe; outside light purplish pink (63D); the reddish style and stamens are prominently exerted; moderately scented. Calyx 2mm long, moderate greenish blue (114B). Lvs obovate, 74 × 36mm, cuneate at base, broadly acute at apex, margins flat, moderate olive green (137A)

above, with colourless hairs predominantly on the veins below. Shrub 2.1. × 1.2m in 9 years.

Mid-season (mid April in central Alabama, USA). Hardy to -16C.

Nomenclatural standard: colour print of an electronic image provided by M.M. Gibson (WSY0125634).

Etymology: named for a volunteer at the Huntsville Madison County Botanical Garden (USA).

Based on proximity of other plants and characteristics of this selection the pollen parent was probably *canescens*. The seed parent was a seedling of a plant grown by John Anderson (deceased – a nurseryman of Scottsboro, Alabama, USA) which he called "best red", from which seed was collected by Vernon Bush in 2006.

(r) **'Drifting Dreams'**

cv. Elepidote rhododendron: 'Summer Sunrise' (s) × 'Plum Passion'  
**H:** J. Barlup (2009), **G:** J. Barlup (2013), **N:** J. Barlup (2016), **REG:** J. Barlup (2018)  
Fls 13/ball truss, broadly funnel-shaped, 51 × 76mm, with 7 rounded, emarginate and slightly wavy-edged lobes, deep purplish pink (54B) in bud, opening inside light purplish pink (65B), shading to pale purplish pink (65D) at the margins, with strong red (46A) nectar pouches (larger on the dorsal side) and strong red (46A) spots below the dorsal lobe; outside light purplish pink (65B), shading to pale purplish pink (65D) at the margins; filaments white, anthers whitish; style white, with a prominent red-pink stigma; not scented. Calyx insignificant. Lvs elliptic, 133 × 54mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A), hairless. Shrub 0.9 × 1.2m in 8 years. Mid-season (mid May in Seattle area, Washington, USA). Hardy to -15C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108643).

(r) **'Ed's Heavenly Sunshine'**

cv. Elepidote rhododendron: 'Janet Blair' (s) × 'Evening Glow' × 'Crest'  
**H:** H.E. Reiley (2007), **G:** M.L. Reiley (2018), **N:** S.L. Reiley (2018), **REG:** Reiley Ridge Nursery (2018)  
Fls 8–10/ball truss, broadly funnel-shaped, 76mm wide, with 5–7 rounded, wavy-edged lobes, strong orange-yellow (22A) in bud, opening inside pale greenish yellow (10D), with light yellow (10B) at throat, midveins light yellow-green (145B) on some flowers, strong purplish red (64B) on others; outside light yellow (10C); stamens very short, perhaps sterile. Calyx 2mm long, strong yellow-green (143C). Lvs oblong, 102 × 38–51mm, rounded at base, broadly acute at apex, with margins downcurved, matt, moderate olive green (146A), hairless. Shrub 1.2 × 1.2m in 10 years. Mid-season (May in Maryland, USA). Hardy to -15C.  
Etymology: named by the hybridiser's daughter and incorporating his name.

(r) 'Emilene'

cv. Elepidote rhododendron: 'Roseum Elegans' × Fabia Group (s) × 'Goldkrone'

**H:** J. Jäck (1996), **G:** J. Jäck (2003), **N:** J. Jäck (2018), **REG:** J. Jäck (2018)

Fls single, 17–19/dense, slightly flattened truss, funnel-shaped, 40–45 × 75–80mm, with 5 rounded and flat-edged lobes, buds darkish pink, opening pale pink, with dark red-pink spotting on the dorsal lobe; outside pale pink, lightly washed with darker tones. Lvs oblanceolate, 90–130 × 40–50mm, obtuse at base, rounded and mucronate at apex, with flat margins, glossy and green above, hairless below. Shrub 1.1 × 1.1m in 22 years. Early May. Hardy to –22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125507). Etymology: named for the wife of a Brazilian friend of the registrant.

(r) 'Emily Fisher'

cv. Elepidote rhododendron: un-named seedling (s) × 'Lem's Cameo'

**H:** C. Mugridge (2002), **G:** C. Mugridge (2012), **N:** C. Mugridge (2018), **I:** Millais Nurseries (2018), **REG:** C. Mugridge (2018)

Fls 8–10/lax truss, broadly funnel-shaped, 50 × 100mm, with 5–6 wavy-edged lobes, strong purplish red (63A) in bud, opening inside light orange-yellow (16B), with a broad red ring around the base of the throat, but without spotting, the lobe sinuses suffused purplish pink; outside light orange-yellow (16B); not scented. Calyx 10–40mm long, pale orange/pink. Lvs elliptic, 150 × 50mm, rounded at base, broadly acute at apex, with a satin-like upper surface, hairless. Shrub 2.0 × 1.5m in 15 years. Mid-season. Hardy to –15C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125635). Etymology: named for the registrant's daughter-in-law.

(r) 'Ercilia'

cv. Elepidote rhododendron: 'Crimson Pippin' (s) × *rex*

**H:** K. Heger (1992), **G:** J. Jäck (2000), **N:** J. Jäck (2015), **REG:** J. Jäck (2018)

Fls single, 18–19/dense, dome-shaped truss, campanulate, 40 × 55mm, with 7 wavy-edged lobes, buds dark rose pink, opening inside dark rose pink, fading to light rose pink, with dark pink-red at base; outside dark rose pink, fading to light rose pink. Lvs lanceolate to oblanceolate 130–170 × 60–65mm, obtuse at base, acute and mucronate at apex, with flat margins, semi-glossy and green above, with a thick, ochre-coloured indumentum below. Shrub 2.2 × 2.0m in 26 years. Late April to early May. Hardy to –22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125509). Etymology: named for the mother of the exchange student of the registrant.

(r) 'Exbury Centenary'

cv. Elepidote rhododendron: *sinogrande* (s) × unknown

**H:** Exbury Gardens, **G:** Exbury Gardens, **N:** T. Clarke (2018), **REG:** T. Clarke (2018)

Fls 23/domed truss, ventricose-campanulate, 60 × 60mm, with 8 wavy-edged, shallowly notched lobes, buds strong purplish red (64C), opening inside light purplish pink (65B), with strong purplish red (63B) at corolla margins, aging to light yellow-green (150D), with light purplish pink (63D) at corolla margins; outside similarly coloured; with a dark red (59A) blotch at the base of the dorsal throat; filaments white, anthers brown; stigma prominent, lime green. Calyx insignificant, light greenish grey (190C). Lvs oblanceolate, 300 × 120mm, oblique at base, obtuse at apex, matt green above, with a yellowish grey (156A) indumentum below when young, which ages to a slightly more fawn colour. Shrub 3–4 × 2m in 10 years. April. Hardy to –7C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125636). Etymology: named to celebrate 100 years of the de Rothschild ownership of Exbury Gardens.

(r) 'Fliedertraum'

cv. Elepidote rhododendron: 'Midnight Mystique' (s) × 'Domino' (Hachmann)

**H:** B. Wickhorst (2004), **G:** Baumschule H.

Hachmann, **N:** B. Wickhorst, **I:** Baumschule H.

Hachmann (2018), **REG:** O. Tschetsch (2018)

Fls 12–15/flattened ball truss, funnel-shaped, 30 × 85mm, with 5 flat lobes, buds vivid reddish purple (74A), opening inside pure white, with 12mm wide margins of light purple (75A), throat and dorsal lobe with a blotch of deep red (60A), surrounded by light yellow (10C) spots, with light reddish purple (74C) at the extreme base; outside light reddish purple (74C); 10 stamens, filaments deep purplish pink (68A), with light purplish pink (68D) at base, anthers coffee brown; style light greenish yellow, stigma olive green, with a dark red margin and a star-shaped pattern of lines; ovary dark olive green, with short white hairs; not scented. Calyx 10mm long, strong brown (172A), with some light yellow-green (144D). Pedicels 40mm long, strong brown (172A) and strong yellow-green (145A). Lvs elliptic, 145 × 65mm, oblique at base, acute at apex, with margins upcurved, moderate olive green (147A) above, strong yellow-green (144B) below, hairless; petioles 20mm long, greyish red (184A). Shrub 0.8 × 1.2m in 10 years. Mid to end May (in northern Germany). Hardy to –22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125588).

(a) **'Galapagos'**

cv. Evergreen azalea: 'Mariko' (s) × 'Blue Danube' (1969)

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2–3/truss, funnel-shaped, 25 × 45mm, with 5 wavy-edged lobes, buds strong reddish purple (70B), opening inside reddish purple (74B–C), with vivid red (45A) spots on the dorsal and adjacent lobes; outside light reddish purple (74C); 7 stamens, filaments vivid reddish purple (74B), with very pale purple (69D) at base, anthers dark red (183A); style strong purplish red (58B), stigma strong yellowish pink (34D). Calyx 2mm long, light yellow-green (145B). Pedicels 11mm long, strong yellow-green (144A/B), with greyish red (178A). Lvs obovate, 33 × 17mm, rounded at base, broadly acute at apex, margins upcurved, glossy olive green, darker than (147A), with white hairs at tip and brown hairs at base above, moderate olive green (146A), with brown hairs below, midrib with longer brown hairs; petioles 3mm long, with long brown hairs. Shrub 0.3 × 0.8m in 10 years. Mid May to early June (in northern Germany). Hardy to –24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125546).

Etyymology: named after the Galapagos National Park in Ecuador.

(r) **'Graziella Blue'**

cv. Elepidote rhododendron: 'Filigran' (s) × 'Anatevka'

**H:** B. Wickhorst (2008), **G:** Baumschule H. Hachmann, **N:** B. Wickhorst, **I:** Baumschule H. Hachmann, **REG:** O. Tschetsch (2018)

Fls 17–20/ball truss, funnel-shaped, 30 × 80mm, with 5 wavy-edged lobes, buds deep reddish purple (77A), opening inside moderate purplish pink (78D), overlaid with strong reddish purple (78A), with merging markings of dark greyish purple (202A) and deep red (185A) on the dorsal and adjacent lobes, nectaries deep purplish red (59B); outside light purple (78C), overlaid with vivid reddish purple (78A); 10 stamens, filaments moderate purplish pink (74D), paling to very pale purple (69D) towards the base and apex, anthers light purple (80D), with touches of pale yellow-green (4D); style varying shades of pink (65A/48B/27D), stigma dark red (59A); ovary deep yellowish green (141B), with short white hairs; not scented. Calyx 3mm long, pale red and very pale green. Pedicels 27mm, strong yellow-green (145A). Lvs linear, 90 × 12mm, cuneate at base, acuminate at apex, with decurved and rugose margins, glossy, moderate olive green (147A) above, strong yellow-green (145A) below, with light yellow-green (145B/C) at midrib, hairless; petioles 6–10mm, moderate reddish brown (177A), dark red (183A) and strong yellow-green (144A). Shrub 1.0 × 1.0m in 10 years. Mid May to early June (in northern Germany). Hardy to –24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125591).

(r) **'Hachmann's Black Eye'**

cv. Elepidote rhododendron: 'Hachmann's Mamamia' (s) × 'Midnight Beauty'

**H:** Holger Hachmann (2009), **G:** Baumschule H. Hachmann, **N:** Holger Hachmann, **I:** Baumschule H. Hachmann, **REG:** O. Tschetsch (2018)

Fls 14–17/ball truss, funnel-shaped, 35 × 70mm, with 6–7 very wavy-edged lobes, buds vivid purple (81A), with deep purple (83B) at base, opening inside vivid reddish purple (74A/B), overlaid with vivid reddish purple (78A) at midrib and edges, with a large blotch, becoming spots, of dark purple (79A/202A) on the dorsal and adjacent lobes; outside vivid reddish purple (78A), with deep purplish red (71A) at midrib; 12–13 stamens, filaments strong purplish red (60B), anthers moderate orange-yellow (165C); style 3mm diameter, strong purplish red (58B), stigma deep red (60A); ovary green, hairless; not scented. Normal calyx absent. Lvs elliptic, 150 × 55mm, cuneate at base, broadly acute at apex, with wavy and upcurved margins, matt, moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 25mm long, greyish brown (166A). Shrub 1.1 × 1.2m in 10 years. Mid to end May (in northern Germany). Hardy to –23C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125594).

(r) **'Hachmann's Eddy'**

cv. Elepidote rhododendron: 'Olga' (s) × 'Norfolk Candy'

**H:** Holger Hachmann (1997), **G:** Baumschule H. Hachmann, **N:** Holger Hachmann, **I:** Baumschule H. Hachmann (2018), **REG:** O. Tschetsch (2018)

Fls 11–16/ball truss, funnel-shaped, 40 × 70mm, with 5–6 wavy-edged lobes, buds blends of pink and yellowish pink (39B/47C/48A), opening inside moderate yellowish pink (31D), with pale orange-yellow (23D) at margins and blends of light orange-yellow (19A) and pale yellow (20C) at the centre, overlaid with vivid orange-yellow (23A) and brilliant yellow (9C), with a moderate reddish brown (175A) blotch and strong reddish orange (169A) spots in the throat and on the dorsal and adjacent lobes; outside strong reddish orange (31B), with pale orange-yellow (23D) margins and strong red (39A) at midrib; 10 stamens, filaments pale yellow (8D/10D), anthers pale yellow (18C); style yellow-green (154C–A), stigma olive green, with a very thin red margin; ovary red-orange, with an olive green tip and short white hairs; not scented. Calyx 10mm long, strong orange (25B), spotted with a deeper tone. Lvs elliptic, 100 × 35mm, rounded at base, broadly acute at apex, with flat margins, glossy, moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles strong yellow-green (144B). Shrub 0.9 × 1.3m in 10 years. Early to late May (in northern Germany). Hardy to –23C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125597).

(r) **'Hachmann's Euphoria'**

cv. Elepidote rhododendron: 'Raphaela' (s) × 'Hachcher' (CHERRY KISS)

**H:** B. Wickhorst (2006), **G:** Baumschule H.

Hachmann, **N:** B. Wickhorst, **I:** Baumschule H.

Hachmann, **REG:** O. Tschetsch (2018)

Fls 12–15/conical truss, funnel-shaped, 50 × 80mm,

with 5 wavy-edged lobes, buds dark red, opening inside pure red, with a prominent, spotted, dark greyish purple (202A) blotch on the dorsal and adjacent lobes; outside pure red, with faint markings in a darker tone; 9–10 stamens, filaments vivid purplish red (66A–B), anthers pale yellow (18C); style vivid purplish red (66B) with green at base, stigma dark greyish purple (202A); not scented.

Calyx 1mm long, moderate reddish brown (177A).

Pedicels 40mm long, strong yellowish green (144A).

Lvs elliptic, 160 × 60mm, rounded at base, acute at apex, with wavy and decurved margins, matt, moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 26mm long. Shrub 1.0 × 1.2m in 10 years. Mid to late May (in northern Germany). Hardy to –24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125599).

(r) **'Hachmann's Melissa'**

cv. Elepidote rhododendron: 'Sariska' (s) × 'Hachpico' (HACHMANN'S PICOBELLO)

**H:** Holger Hachmann (2004), **G:** Baumschule H.

Hachmann, **N:** Holger Hachmann (2018),

**I:** Baumschule H. Hachmann, **REG:** O. Tschetsch (2018)

Fls 12–14/domed truss, funnel-shaped, 30 × 80mm,

with 5 wavy-edged lobes, buds light purplish pink (62C), opening inside pale purplish pink (65D), with a large blotch of dark red (187A), surrounded by paler red spots (60A–C) on the dorsal and adjacent lobes; outside pale purplish pink (65C–D), with moderate purplish pink (65A) veins; 12–13 stamens, filaments yellowish white (155D), anthers moderate orange-yellow (164B); style brilliant yellow-green (150C), with strong yellow-green (144B) at base, stigma brilliant yellow-green (154B); ovary strong yellow-green (144B), with short white hairs. Calyx 3mm long, light yellow-green (145D). Pedicels 40mm long, strong yellow-green (145A). Lvs elliptic,

135 × 50mm, rounded at base, broadly acute at apex, with margins decurved, matt, opening strong yellow-green (144A), with short white hairs, becoming moderate olive green (147A) to dark yellowish green (139A) above, moderate yellow-green (146C) below, hairless; petioles 20mm long, light yellow-green (144D). Shrub 1.2 × 1.5m in 10 years. Early to late May (in northern Germany). Hardy to –22C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125603).

(r) **'Harold Greer'**

cv. Elepidote rhododendron: 'Very Berry' (s) × 'Horizon Monarch'

**H:** H.E. Greer (1997), **G:** H.E. Greer (2012),

**N:** D. Manion (2017), **REG:** D. Manion (2018)

Fls 10/ball truss, open funnel-shaped, 47 × 90mm,

with 5 rounded, very wavy-edged lobes, deep purplish pink (55A) in bud, opening inside pale yellow (18C), with deep purplish pink picotee margins, and with nectar pouches and small dorsal flare at base of strong purplish red (54A); outside strong purplish red (54A); filaments and style whitish. Calyx 13–32mm long, pale yellow (11C), with deep purplish pink (55A) stripes. Lvs elliptic, 152 × 70mm, rounded at base, acute at apex, margins slightly upcurved, matt and moderate olive green (146A) above, hairless. Shrub 1.5 × 1.5m in 15 years. Mid-season (early May in western Oregon, USA). Hardy to –18C.

Nomenclatural standard: colour print of an electronic image provided by the hybridiser (WSY0125640).

(r) **'Harwood Dale'**

cv. Elepidote rhododendron: un-named seedling (s) × 'Old Port'

**H:** C. Mugridge (2000), **G:** C. Mugridge (2016),

**N:** C. Mugridge (2018), **I:** Millais Nurseries (2018),

**REG:** C. Mugridge (2018)

Fls 10–13/ball truss, broadly funnel-shaped,

35 × 70mm, with 5 wavy-edged lobes, buds deep mauve, opening inside light purple (75A), with strong purplish red (61B) blotch and spots on the dorsal and adjacent lobes; outside very pale purple (75D); filaments and anthers white; style white, with a reddish pink stigma; not scented. Calyx 1mm long, pale green. Lvs elliptic, 130 × 50mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy above, hairless. Shrub 1.0 × 1.0m in 18 years.

Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125560).

Etymology: named after the site on the North Yorkshire moors where the ashes of the registrant's parents were scattered.

(r) **'Helene Richter'**

cv. Elepidote rhododendron: *makinoi* (s) × *decorum*

**H:** H.L. Meltzer (1988), **G:** H.L. Meltzer (2001),

**N:** H.L. Meltzer (2005), **I:** H.L. Meltzer (2005),

**REG:** H.L. Meltzer (2018)

Fls 9–11/truss, open funnel-shaped, 70–80mm wide, with 5–6 slightly wavy-edged lobes, rose pink in bud fading as it comes out, the fully developed flower almost pure white, with an inconspicuous group of yellowish spots at the base of the dorsal throat, slightly pink tinged on the outside. Lvs narrowly elliptic, cuneate at base, acute to shortly acuminate at apex, margins flat, glossy green above. Shrub 0.4 × 0.6m in 10 years. Mid to late May. Hardy to –27C.

Nomenclatural standard: colour print provided by the registrant (WSY0108604). Has been seen as "Macdec540". The parental cross was referred to as 15/1988.

(r) 'Hesse's Daydreamer'

cv. Elepidote rhododendron: 'Hachkaba' (HACHMANN'S KABARETT) (s) × 'Trianon' (H: H. Hesse (2000), G: H. Hesse (2006), N: H. Hesse, REG: O. Tschetsch (2018))  
Fls single, 14/ball-shaped truss, funnel-shaped, 45 × 100mm, with 7 wavy-edged lobes, inside very pale purple (69D), with touches of a deeper tone (69C), throat washed with light to pale yellow (11B–C), with merging dark red (187A–B) and dark purplish red (59A–B) spotting on the dorsal and adjacent lobes; outside very pale purple (69D), with deeper tones (69A–B) at midribs; 14 stamens, filaments pale yellow (8D), with strong purplish red (60D) at base, anthers light yellow (160B); style brilliant yellow-green (150C), with vivid red (44A) at base, stigma brilliant yellow-green (150A); ovary moderate yellow-green (137C), with short white hairs; not scented. Calyx 1mm long, strong yellow-green (144C). Pedicels strong yellow-green (144C/144B). Lvs elliptic, 160 × 70mm, oblique at base, broadly acute at apex, margins wavy, moderate olive green (147A), with strong yellow-green (144C) at midrib above, moderate yellow-green (146B), with strong yellow-green (145A) midrib below, with a few white hairs when young, quickly becoming hairless; young shoots dark red (187A). Shrub 1.1 × 1.4m in 18 years. Late May to early June (in northwest Germany). Hardy to –22C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125538).

(r) 'Hesse's Leonie'

cv. Elepidote rhododendron: 'Consolini's Windmill' (s) × 'Hachmann's Berolina'  
H: H. Hesse (2002), G: H. Hesse (2008), N: H. Hesse, REG: O. Tschetsch (2018)  
Fls single, 8–12/domed truss, funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, inside pale yellow-green (155A), with a 6mm wide margin of purplish pink (66A–C), base of the dorsal and adjacent lobes light yellow-green (150D), overlaid with strong yellow-green (144B–C) spots, lower lobes spotted on margins at base, with very pale midribs; outside purplish pink (66A–C), with moderate purplish red (70A) at base and vivid red (57A) along the midribs; 7 stamens, filaments greenish white (157D), overlaid with light purplish pink (62C), anthers moderate reddish brown (175A); style translucent, moderate purplish red (59C), with pale yellow-green (4D) at base, stigma dark red (187A). Calyx 6mm long, light yellow-green (145C), overlaid with vivid purplish red (57B) at base. Pedicels 30mm long, strong yellow-green (144C), with vivid purplish red (57B) stripes. Lvs elliptic, 120 × 45mm, oblique at base, broadly acute at apex, margins twisted, matt, dark green, hairless; petioles 25mm long, moderate yellow-green (146B), with greyish red (176A) above, strong yellow-green (144C) below. Shrub 1.2 × 1.4m in 16 years. Late May to early June (in northwest Germany). Hardy to –22C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125542).

(r) 'Hotel Paradisco'

cv. Elepidote rhododendron: parentage unknown  
H: USA origin (pre–1992), G: J. Jäck (2000), N: J. Jäck (2000), REG: J. Jäck (2018)  
Fls single, 5–8/loose, open-topped truss, funnel-shaped, 60 × 80–90mm, with 7 emarginate, wavy-edged lobes, buds dark yellow-pink, opening inside pale pink, with creamy yellow tones towards base, becoming very pale pink, shading through pale creamy yellow to bright greenish yellow at base; outside pale pink, with creamy yellow towards base; very fragrant. Lvs oblanceolate, 140–170 × 45–60mm, semi-glossy and green above, hairless. Shrub 2.5 × 1.5m in 25 years. Early to mid May. Hardy to –22C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125513).  
Etymology: named after a hotel on Mallorca.

(a) 'Iguazu'

cv. Evergreen azalea: 'Mariko' (s) × 'Blue Danube' (1969)  
H: F. Schnupper (2003), G: F. Schnupper, N: F. Schnupper, I: Schnupper Baumschule (2018), REG: O. Tschetsch (2018)  
Fls single, 2–4/truss, funnel-shaped, 30 × 50mm, with 5 wavy-edged lobes, buds vivid reddish purple (74B), ribbed with moderate purplish pink (68C), opening inside light reddish purple (74C), with vivid reddish purple (74A) spots, merging with vivid reddish purple (66A) spots on the dorsal and adjacent lobes; outside vivid reddish purple (74B–C); 7–8 stamens, filaments deep purplish pink (66C) and very pale purple (69D); style vivid purplish red (66A), with deep purplish pink (66C) at base. Calyx 2mm long, strong yellow-green (144C), with white hairs. Pedicels 12mm long, strong yellow-green (144B), marked with some strong brown (172A) and greyish red (184A) and with white hairs. Lvs obovate, 40 × 16mm, rounded at base and apex, with margins upcurved, glossy, moderate olive green (147A), with a margin of light brown hairs above, moderate yellow-green (147C), with sparse brown hairs below; petioles 5mm long. Shrub 0.4 × 0.8m in 10 years. Mid May to early June (in northern Germany). Hardy to –24C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125547).  
Etymology: named after the Iguazú National Park in Argentina.

(r) 'Jack's Choice'

cv. Elepidote rhododendron: parentage unknown  
H: R.A. Furman (c.2002), G: R.A. Furman (2010), S: K.F. Humphries (2017), N: K.F. Humphries (2018), REG: K.F. Humphries (2018)  
Fls 16–18/ball truss, funnel-shaped, 64 × 90mm, with 7–8 rounded, overlapping, wavy-edged lobes, buds strong purplish red (61B), opening inside moderate purplish pink (66D), shading to pale orange-yellow (159C) at base, with short rays of dark red (183B) at throat, the longest at the base of the dorsal lobe; outside deep purplish pink (66C); stamens absent; style orange-yellow, with brown stigma; lightly scented. Calyx 19mm long, pale

orange-yellow (159B). Lvs elliptic, 127 × 64mm, rounded at base, obtuse at apex, margins wavy and up-angled from the midvein, semi-glossy, moderate olive green (147A), hairless. Shrub 2.1 × 2.1m in c.15 years. Mid-season (mid May on Cape Cod, Massachusetts, USA). Hardy to -21C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125568).  
Etymology: named for Jack Donahue, husband of the registrant and son-in-law of the hybridizer.

(r) 'Jaerperlå'

cv. Elepidote rhododendron: 'Humboldt' (s) × *wardii* (Ludlow & Sherriff form)  
**H:** O. Fosse (2005), **G:** O. Fosse (2009), **N:** O. Fosse (2009), **I:** O. Fosse (2015), **REG:** P.M. Jørgensen (2018)  
Fls 5–12/open truss, open funnel-shaped, 40–50 × 20–40mm, with 5 more-or-less flat-edged lobes, violet in bud, opening inside whitish, pale violet outside; with five deep burgundy basal blotches around the base of the throat, largest on the dorsal side, where they are surmounted by some dorsal spotting (on a slightly yellowish ground). Calyx 1–2mm long, green. Lvs lanceolate, 60–120 × 30–40mm, rounded at base, pointed at apex, margins flat, very shiny green above, hairless. Shrub 1.5 × 1.0m in 10 years. May. Hardy to -30C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125641). Etymology: the epithet translates from Norwegian as "Pearl of Jåren", Jåren being a district in western Norway. Published in *Lapprosen* 29 (1): 24 (2018).

(r) 'Jen Skerman'

cv. Elepidote rhododendron: 'Jeanette Clarke' (s) × *arboreum*  
**H:** L. Begg (2003), **G:** J.C. Gray (2010), **N:** J.C. Gray (2018), **I:** Brindabella Country Gardens (2018), **REG:** J.C. Gray (2018)  
Fls 14/ball truss, campanulate, 60 × 80mm, with 5 wavy-edged, shallowly crenate lobes, buds strong red (50A), opening inside deep purplish pink (55A), with a stripe of strong pink (48D) at midrib, lightly spotted red on dorsal lobe; outside deep purplish pink (55A). Lvs elliptic, 150 × 45mm, acute at base and apex, with flat margins, glossy above, hairless. Shrub 2.0 × 2.0m in 10 years. September (in Queensland, Australia). Very heat tolerant.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125570).

(r) 'John Anderson'

cv. Elepidote rhododendron: *arizelum* (s) × unknown  
**H:** Exbury Gardens, **G:** Exbury Gardens, **N:** T. Clarke (2018), **REG:** T. Clarke (2018)  
Fls 31/conical truss, ventricose-campanulate, 55 × 67mm, with 8 wavy-edged lobes, inside pale greenish yellow (1D); outside light greenish yellow (1C); with a dark red (187A) blotch at the base of the throat (covering the three dorsal nectar pouches) and extending for 10mm above; filaments white, anthers dark reddish brown; prominent lemon yellow style and stigma. Calyx insignificant, pale yellow-green

(193D). Lvs oblanceolate, 250 × 102mm, obliquely cuneate to truncate at base, acute at apex, margins wavy, with a brownish orange (164A) indumentum below at first, which ages to moderate orange-yellow (164B). Small, multi-stemmed tree, 8.0 × 6.0m in 30 years. April. Hardy to -7C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125642).  
Etymology: named for a former head gardener at Exbury Gardens.

(r) 'Kailey'

cv. Elepidote rhododendron: 'Polarnacht' (s) × 'Wild Berry'  
**H:** J. Barlup (2009), **G:** J. Barlup (2014), **N:** J. Barlup (2016), **REG:** J. Barlup (2018)  
Fls 13/ball truss, broadly funnel-shaped, 51 × 76mm, with 5 rounded, very wavy-edged lobes, buds deep purplish red (71A), opening inside strong purple (84A), with light purple (84B) midveins on all lobes and with a dark greyish reddish brown (200A) blotch on the dorsal lobe; outside strong purple (84A), with light purple (84B) midveins on all lobes. Calyx insignificant. Lvs elliptic, 114 × 51mm, cuneate at base, broadly acute at apex, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 8 years. Mid-season (mid May in Seattle area, Washington, USA). Hardy to -18C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108646).  
Etymology: named for a business associate of the hybridiser.

(a) 'Kakadu'

cv. Evergreen azalea: 'Mariko' (s) × 'Blue Danube' (1969)  
**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)  
Fls single, 2–3/truss, funnel-shaped, 20 × 40mm, with 5 wavy-edged lobes, buds moderate purplish red (70A) and strong reddish purple (70B), with pale purplish pink (56C) ribs, opening inside vivid reddish purple (78A), with vivid purplish red (66B) spots on the dorsal and adjacent lobes; outside strong reddish purple (78B), with strong purplish red (61B) ribs; 7 stamens, filaments vivid reddish purple (74A) and very pale purple (75D), anthers dark red (187A); style strong purplish red (54A), with brownish orange (165B) at base. Calyx 2mm long, strong yellow-green (144B). Pedicels 10mm long, strong yellow-green (144B), with some deep red (185A). Lvs obovate, 30 × 16mm, rounded at base and apex, with margins upcurved, glossy, moderate olive green (147A) above, moderate yellow-green (146B), with sparse white hairs, except for midrib which has brown hairs, below; petioles 5mm long. Shrub 0.4 × 0.8m in 10 years. Mid May to early June (in northern Germany). Hardy to -24C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125548).  
Etymology: named after the Kakadu National Park in Australia.

(a) **'Komodo'**

cv. Evergreen azalea: 'Mariko' (s) × 'Satschiko'

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2–3/truss, funnel-shaped, 30 × 40mm, with 5 wavy-edged lobes, buds deep pink to strong pink (50B–50C), opening inside deep pink (52B–C), with strong red (53C) spots on the dorsal and adjacent lobes; outside deep pink (52B–C); 8 stamens, filaments deep red (53A), anthers dark greyish purple (202A); style strong purplish red (58B), stigma moderate orange (167A). Calyx 2mm long, light yellow-green (145B). Pedicels 8mm long, strong yellow-green (144C), with moderate red (180A). Lvs elliptic, 26 × 16mm, cuneate at base, broadly acute at apex, with flat margins, glossy, moderate olive green (146A) above, moderate yellow-green (146C), with white hairs, except along midrib, which has brown hairs, below; petioles 5mm long, strong yellow-green (144C). Shrub 0.4 × 0.7m in 10 years. Mid to end May (in northern Germany). Hardy to –24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125549).  
Etymology: named after the Komodo National Park in Indonesia.

(a) **'Kurzeme'**

cv. Deciduous azalea: 'Homebush' (s) × 'Austra'

**H:** R. Kondratovičs (1999), **G:** R. Kondratovičs (2006), **N:** G. Riekstiņa (2018), **REG:** Rhododendron Nursery Babīte, University of Latvia (2018)

Fls double, 15/strongly domed truss, tubular funnel-shaped, 35 × 50mm, with 2 whorls of 5 flat lobes, buds vivid reddish orange (44B), opening inside deep pink (52B), with pale purplish pink (56C) at midrib; outside pale purplish pink (56D); style pink, stigma green; lightly scented. Calyx 2mm long, green. Lvs elliptic, 80 × 35mm, acute at base and apex, glossy above, with small, white hairs along the wavy margins. Shrub 2.3 × 2.3m in 18 years. Late May to early June. Hardy to –27C.

Nomenclatural standard: colour print provided by the registrant (WSY0125572). Etymology: named after one of the Latvian historical and cultural regions.

(r) **'Ladylike'**

Syn. of 'Lady Like'.

(r) **'Lady Like'**

cv. Elepidote rhododendron: 'Hachraba' (RABATZ) (s) × 'Domino' (Hachmann)

**H:** Holger Hachmann, **G:** Baumschule H. Hachmann, **N:** Y. Dibaba (2018), **I:** Vorwerk Pflanzenhandel GmbH (2018), **REG:** O. Tschetsch (2018)

Fls 10–13/ball truss, funnel-shaped, 35 × 95mm, with 5 wavy to frilly-edged lobes, buds vivid reddish purple (74A), opening inside purple (74B–75C), with a large dark red (187A) blotch, changing to dark red (59A) spots on the three dorsal lobes, with a lighter covering of dark red (59A) spots on the ventral lobes; outside purple blends (74B–D/75C); 10 stamens, filaments yellowish white (155D),

anthers moderate yellow (161B); style light yellow-green (150D), stigma dark red (183B). Calyx 1mm long, pale yellow (165D). Pedicels 45mm long, strong yellow-green (144C). Lvs elliptic, 145 × 55mm, oblique at base, broadly acute at apex, with margins wavy and twisted, glossy, opening moderate yellow-green (146B), with short white sticky hairs, becoming moderate olive green (147A) above, moderate yellow-green (146B) below, hairless; petioles 30mm long, strong yellow-green (144B). Shrub 1.1 × 1.5m in 10 years. Early to late May (in northern Germany). Hardy to –22C.

Syn: 'Ladylike'

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125605).

(a) **'Latgale'**

cv. Deciduous azalea: 'Homebush' (s) × 'Austra'

**H:** R. Kondratovičs (1999), **G:** R. Kondratovičs (2006), **N:** G. Riekstiņa (2018), **REG:** Rhododendron Nursery Babīte, University of Latvia (2018)

Fls double, 14/slightly domed truss, tubular funnel-shaped, 40 × 45mm, with 2 whorls of 5 flat lobes, buds strong yellowish pink (32C), opening inside pale purplish pink (56A), with a lighter tone (56D) at midrib; outside pink, with yellowish white (N155D) at midrib; style pink, stigma green; lightly scented. Calyx 2mm long, green. Lvs elliptic, 90 × 35mm, acute at base and apex, glossy above, hairless. Shrub 2.6 × 2.6m in 18 years. Late May to early June. Hardy to –27C.

Nomenclatural standard: colour print provided by the registrant (WSY0125573). Etymology: named after one of the Latvian historical and cultural regions.

(r) **'Latvijai 100'**

cv. Elepidote rhododendron: 'Falling Snow' (s) × 'Erato'

**H:** G. Riekstiņa (2006), **G:** G. Riekstiņa (2013), **N:** G. Riekstiņa (2018), **REG:** Rhododendron Nursery Babīte, University of Latvia (2018)

Fls 18/conical truss, open funnel-shaped, 45 × 70mm, with 5 strongly frilled lobes, buds deep red (60A), opening inside very pale purple (75D) at throat and centre of lobes, with broad, strong purplish red (61B), margins, dorsal and adjoining lobes spotted with strong yellowish green (N144A); outside strong purplish red (61B); filaments light pink, with white anthers; style light red, with dark cherry red stigma. Calyx 1mm long, green. Lvs elliptic, 140 × 60mm, rounded at base, acute at apex, with flat margins, glossy above, hairless. Shrub 1.0 × 1.5m in 11 years. Hardy to –26C.

Nomenclatural standard: colour print provided by the registrant (WSY0125574). Etymology: named in honour of one hundred years since Latvia became an independent state in 1918.

(r) **'Liquid Sunshine'**

cv. *Elepidote rhododendron*: 'Lemon Prelude' (s) × 'Goldprinz'

**H:** J. Barlup (2005), **G:** J. Barlup (2009),

**S:** R. Blackmore (2017), **N:** R. Blackmore (2017),

**REG:** R. Blackmore (2018)

Fls 17/domed truss, broadly funnel-shaped to saucer-shaped, 44 × 83mm, with 7 rounded, emarginate or pinched lobes, with wavy-edged margins, strong red (39A) in bud, opening inside pale yellow (11C), with 3–4 strong red (53B) spotted flares extending 19mm from the throat into the dorsal area; outside pale yellow (11C); pistil and stamens yellow. Calyx 19mm long, pale yellow (11C), with strong red (53B) spots. Lvs elliptic to oblanceolate, 95 × 44mm, rounded to slightly cuneate at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.8m in 6 years. Mid-season (May in Victoria, British Columbia, Canada). Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108645).

(r) **'Lois Norma'**

cv. *Elepidote rhododendron*: 'Helen Everitt' (s) × 'Phipps Yellow'

**H:** R. Murcott (1983), **G:** M. Borsky (1987),

**N:** M. Borsky (1993), **REG:** M. Borsky (2018)

Fls 9–10/ball truss, funnel-shaped, 45 × 86mm, with 7 rounded, wavy-edged lobes, strong purplish pink (63C) in bud, opening inside very light purple (75C) at margins, shading through very pale purple (75D) to white (155C), with a small, light yellow-green (145B) flare and stippling shading to strong yellow-green (145A) in the throat; outside shading from very light purple (75C) at margins, through very pale purple (75D), then white (155C), to very light purple (75C) 10mm from the calyx; stamens absent or only vestigial and 8mm long; style yellow-white, with a yellow-green stigma. Calyx 1mm long, strong yellow-green (145A). Lvs elliptic, 142 × 60mm, rounded at base, broadly acute at apex, margins flat, matt and moderate olive green (137A) above, hairless. Shrub 1.75 × 4.0m in 34 years. Mid-season (May in central New Jersey, USA). Hardy to –18C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108648).

Etyymology: named for the wife of the registrant.

(r) **'Makdec'**

cv. *Elepidote rhododendron*: *makinoi* (s) × *decorum*

**H:** H.L. Meltzer (1988), **G:** H.L. Meltzer (1998),

**N:** H.L. Meltzer (2005), **I:** H.L. Meltzer (2005),

**REG:** H.L. Meltzer (2018)

Fls probably up to c.10/hemispherical truss, funnel-shaped, 90mm wide, with 5 very wavy-edged lobes, rose pink in bud, opening white in centre, shading to paler pink at margins; outside rose pink, with a limited amount of orange dorsal spotting. Lvs similar to *makinoi* but larger (to at least 170 × 38mm), with decurved margins, matt green above, with a light brown indumentum below. Shrub 2.9 × 1.6m in 29 years. Mid May. Hardy to –27C.

Nomenclatural standard: colour print provided by the registrant (WSY0108601). Has been seen as "Macdec438". The parental cross was referred to as 15/1988.

(r) **'Malpelo'**

cv. *Elepidote rhododendron*: *yakushimanum* (s) × *hyperythrum*

**H:** F. Schnupper (2001), **G:** F. Schnupper,

**N:** F. Schnupper, **I:** Schnupper Baumschule (2018),

**REG:** O. Tschetsch (2018)

Fls single, 9–12/ball-shaped truss, broadly funnel-shaped, 35 × 70mm, with 5 flat lobes, inside pale purplish pink (62D), with inconspicuous brilliant yellow-green (150C) markings on the dorsal and adjacent lobes; outside pale purplish pink (62D), with moderate purplish pink (62B) along midribs; 10 stamens, filaments white, anthers pale yellow (158A); style yellowish white (155D), stigma pale greenish yellow (160C); ovary green, with sparse white hairs. Calyx inconspicuous. Pedicels 50mm long, light yellow-green (145C). Lvs oblong, 150 × 40mm, rounded at base and apex, with decurved and rugose margins, glossy moderate olive green (147A) above, with moderate orange-yellow (164B) hairs below; petioles 30mm long, green, felted grey. Shrub 0.8 × 1.0m in 10 years. Early to late May (in northern Germany). Hardy to –25C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125550).

Etyymology: named after the Malpelo Marine Nature Reserve in the East Pacific Ocean.

(r) **'Margaret Bates'**

cv. *Elepidote rhododendron*: 'Faggetter's Favourite' (s) × 'Lem's Cameo'

**H:** C. Mugridge (2003), **G:** C. Mugridge (2017),

**N:** C. Mugridge (2018), **I:** Millais Nurseries (2018),

**REG:** C. Mugridge (2018)

Fls 18–22/tall domed truss, broadly funnel-shaped, 50 × 90mm, with 5 slightly frilled lobes, buds deep pink, opening inside light purplish pink (73C), with strong purplish red (63B) spots on the dorsal lobe; outside light purplish pink (73C), with touches of strong purplish pink (68B) at margins; anthers brown; pistil pale green; slightly scented. Calyx 1–2mm long, green. Lvs elliptic, 200 × 65mm, rounded at base, broadly acute at apex, margins flat, matt above, hairless. Shrub 1.0 × 1.0m in 15 years. Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125561).

Etyymology: named for the wife of the registrant.

(r) **'Mary Abair'**

cv. *Elepidote rhododendron*: 'Golden Mist' (s) × 'Mac's Orange Crush'

**H:** J. Barlup (2010), **G:** J. Barlup (2016), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 15/ball truss, broadly funnel-shaped, 51 × 76mm, with 6 rounded, wavy-edged lobes, strong red (39A) in bud, opening inside light greenish yellow (8C) at base, shading to pale yellow (8D), on dorsal lobe(s) shading to light pink (39D), and with twin, strong

greenish yellow (153B) basal flares below the dorsal lobe; outside light greenish yellow (8C) at base, shading to pale yellow (8D), shading to light pink (39D) on the dorsal margin(s); filaments yellow, anthers brown; style yellow, with a reddish stigma. Calyx 22mm long, light greenish yellow (8C), shading to pale yellow (8D), with strong greenish yellow (153B) spots. Lvs elliptic, 146 × 54mm, rounded at base, broadly acute at apex, margins flat, matt and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 7 years. Late (mid June in Seattle region, Washington, USA). Hardy to -18C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108647). Etymology: named for the sister of the registrant.

(a) **'Mary's House'**

cv. Evergreen azalea: parentage unknown  
**H:** W.L. Clagett (c.1998), **G:** W.L. Clagett (2007), **N:** W.L. Clagett (2007), **I:** White's Nursery (2015), **REG:** W.L. Clagett (2018)  
Fls double, 3/terminal shoot, saucer-shaped, 38 × 51mm, with 15–18 rounded, very wavy-edged lobes, strong purplish pink (62A) in bud, opening inside moderate purplish pink (62B) in centre of lobes, shading to pale purplish pink (62D) at margins, and with a bright green throat; outside strong purplish pink (62A); stamens absent; stigma bright orange-red. Normal calyx absent. Lvs elliptic, 38 × 13mm, cuneate at base, acute at apex, margins slightly upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.5 × 0.4m in 10 years. Late (late May in Maryland, USA). Hardy to -21C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125649). Etymology: incorporates the name of the registrant's daughter.

(r) **'Masaya'**

cv. Elepidote rhododendron: 'Lem's Monarch' (s) × *fortunei*  
**H:** F. Schnupper (2002), **G:** F. Schnupper, **N:** F. Schnupper (2018), **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)  
Fls single, 10–16/domed truss, funnel-shaped, 50 × 100mm, with 5–6 wavy-edged lobes, buds strong purplish red (60B), opening inside yellowish white (155D), overlaid at margins with strong purplish pink (68B), the dorsal lobes with merging markings of dark reddish orange (175B) and moderate yellow (162B) at base; outside very pale purple (69C), overlaid with strong purplish red (60B); 12–14 stamens, filaments greenish white (157D), with white hairs at base, anthers moderate orange-yellow (165C); style strong yellow-green (145A), covered with short, white glandular hairs from base, paling to light yellow-green (145C) at top, stigma strong yellow-green (144B), with some red dots; ovary moderate olive green (137A), with short, white glandular hairs; slightly scented. Calyx 5mm long, moderate red (180B). Pedicels 35mm long, strong yellow-green (144B), with dark red (187C). Lvs elliptic, 145 × 50mm, rounded at base, broadly acute at apex, with decurved margins, matt, moderate

olive green (147A) above, strong yellow-green (144A) below, hairless; petioles 30mm long, strong yellow-green (144C), with greyish red (176A). Shrub 0.9 × 1.0m in 9 years. Mid May to early June (in northern Germany). Hardy to -24C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125551). Etymology: named after the Masaya Volcano National Park in Nicaragua.

(r) **'Mike Peterson'**

cv. Elepidote rhododendron: 'Evening Embers' (s) × 'Holli's Hope'  
**H:** J. Barlup (2009), **G:** J. Barlup (2015), **N:** J. Barlup (2016), **REG:** J. Barlup (2018)  
Fls 17/ball truss, broadly funnel-shaped, 44 × 64mm, with 5 rounded, very wavy-edged lobes, deep purplish red (71A) in bud, opening inside very pale purple (76D), shading to strong purple (77B) at margins, with twin dorsal flares of light yellow-green (154D) spots; outside very pale purple (76D), shading to strong purple (77B) at margins; filaments whitish, anthers buff; style pinkish, with a reddish stigma. Calyx insignificant. Lvs elliptic, 102 × 48mm, rounded at base, broadly acute at apex, with decurved margins, matt and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.6m in 8 years. Late (late May in Seattle area, Washington, USA). Hardy to -15C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108650). Etymology: named for a longtime friend of the registrant.

(r) **'Milk and Coffee'**

cv. Elepidote rhododendron: 'Hachgold' (GOLDEN EVEREST) (s) × 'Midnight Beauty'  
**H:** B. Wickhorst (2004), **G:** Baumschule H. Hachmann, **N:** B. Wickhorst, **I:** Baumschule H. Hachmann (2016), **REG:** O. Tschetsch (2018)  
Fls 7–10/domed truss, funnel-shaped, 35 × 65mm, with 5 very wavy-edged lobes, buds strong red (53C), opening inside pale yellow (11C), shading to pale orange-yellow (19D) towards margins, overlaid with strong pink (54C) at margins, deepening to vivid purplish red (57C) at edges, with a dark red (59A) blotch and spots on the dorsal and adjacent lobes; outside pale orange-yellow (19D), with dark red (187B) and deep red (53A) markings; 9–12 (usually 10) stamens; filaments pale orange-yellow (19D), with touches of dark red (187B), anthers brown; style orange, yellow and green, stigma brown-red; ovary light green, with white hairs. Normal calyx absent. Pedicels 25mm long, strong red (46A). Lvs elliptic, 90 × 35mm, cuneate at base, acute at apex, with upcurved margins, matt, moderate olive green (147A) above, moderate yellow-green (146B) below, hairless; petioles 20mm long, dark red (187A). Shrub 1.4 × 1.2m in 10 years. Mid to late May (in northern Germany). Hardy to -22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125609). Etymology: flower colour is reminiscent of café au lait.

(r) **'Moe's Journey'**

cv. *Elepidote rhododendron*: 'Mrs Furnivall' × 'Coronation Day' (s) × 'Elegant Touch'

**H:** J. Barlup (1992), **G:** J. Barlup (1999), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 15/domed truss, broadly funnel-shaped, 51 × 76mm, with 5 rounded, flat or slightly cupped margins, strong red (53C) in bud, opening inside pale yellowish pink (36D), shading to light pink (39D) at margins, and with moderate yellowish pink (39C) midveins; dark red (187B) in the throat, continuing as 22mm twin dorsal flares, extending along the side margins of the dorsal lobe, and with 3 shorter ventral flares (also dark red, 187B); outside pale yellowish pink (36D), shading to light pink (39D) at margins, and with light pink (39D) midveins; pistil prominent, style whitish, shading to pink, and with a reddish stigma. Calyx insignificant. Lvs elliptic, 114 × 51mm, rounded at base, broadly acute at apex, up-angled from the midvein but with flat margins, moderate olive green (147A), hairless. Shrub 0.8 × 0.8 in 6 years. Mid-season (mid May in Seattle area, Washington, USA). Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108649).

Etymology: named for the late Moe Massa, a special rhododendron friend of the registrant from Victoria, British Columbia, Canada).

(r) **'Mr Carl'**

cv. *Elepidote rhododendron*: 'The Honourable Jean Marie de Montague' (s) × 'Babar'

**H:** L.M. Player (1998), **G:** L.M. Player, **N:** L.M. Player, **REG:** L.M. Player (2018)

Fls single, 13/domed truss, funnel-shaped, 76 × 90mm, with 5 wavy-edged lobes, strong red (46A) in bud, opening inside strong red (53D), fading to strong purplish red (54A), with deep red (53A) dorsal spotting originating at base; outside opening strong red (53D), fading to strong purplish red (54A); filaments, style and stigma whitish. Calyx negligible. Lvs elliptic, 152 × 64mm, rounded at base, broadly acute at apex, margins flat, semi-glossy, moderate olive green (147A) above, moderate yellow-green (146B) below, hairless. Shrub 1.2 × 1.5m in 10 years. Mid-season (mid April in eastern North Carolina, USA). Hardy to -21C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125500).

Etymology: named for the father of the registrant.

(r) **'Mrs Daniels'**

cv. *Elepidote rhododendron*: 'Mrs Sam' (s) × *hyperythrum*

**H:** L.M. Player (1998), **G:** L.M. Player (c. 2002), **N:** L.M. Player (post-2002), **REG:** L.M. Player (2018)

Fls single, 12/domed truss, funnel-shaped, 51 × 76mm, the 5 rounded lobes, with flat margins, light purplish pink (55C) in bud, opening inside white, with a strong red (46A) spotted blotch on dorsal lobe; outside white; filaments and style white, stigma pinkish red. Calyx negligible. Lvs elliptic, 102 × 38mm, rounded at base, broadly acute at apex, margins flat, glossy, moderate olive green (137A) above, moderate yellow-green

(138B) below, hairless. Shrub 2.4 × 2.7m in 20 years.

Mid-season (mid April in eastern North Carolina).

Hardy to -21C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125501).

Etymology: named for an aunt of the hybridiser.

(r) **'Mrs Sylvia E. Gray'**

cv. *Elepidote rhododendron*: 'Jeanette Clarke' (s) × *arboreum*

**H:** L. Begg (2003), **G:** J.C. Gray (2010), **N:** J.C. Gray (2018), **I:** Brindabella Country Gardens (2018),

**REG:** J.C. Gray (2018)

Fls 14/ball truss, campanulate, 48 × 55mm, with 5 wavy-edged lobes, deep purplish pink (55A) in bud, opening inside deep purplish pink (54B); outside strong pink (54C); with red spots on the dorsal lobe. Lvs elliptic, 135 × 55mm, matt green, hairless. Shrub 2.0 × 1.5m in 10 years. September (Australia). Very heat tolerant. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108652).

(r) **'Ngorongoro'**

cv. *Elepidote rhododendron*: *fortunei* (s) × 'Ring of Fire' (Thompson)

**H:** F. Schnupper (2004), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 8-9/domed truss, funnel-shaped, 40 × 75mm, with 6 wavy-edged and emarginate lobes, buds strong red (51A), opening inside light yellow (10C), with wide margins of deep pink (52C) and with brilliant greenish yellow (1B) in the throat; outside deep purplish pink (55A), paling to light yellowish pink (159A) at base; 11-15 stamens, filaments light yellow-green (150D), anthers moderate yellow (161A); style strong yellow-green (145A), stigma strong yellow-green (144B); ovary moderate yellowish green (138A), with sparse white hairs; slightly scented. Calyx 3mm long, greyish red (178A), with green at base. Pedicels 30mm long, greyish red (178A) with green at base. Lvs oblong, 80 × 30mm, rounded at base, broadly acute at apex, with wavy and decurved margins, strong yellow-green (144A) above, strong yellow-green (144C), with strong greenish yellow (153C) midrib below, hairless; petioles 17mm long, strong orange-yellow (163B). Early to late May (in northern Germany). Hardy to -24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125552).

Etymology: named after the Ngorongoro Conservation Area in Tanzania.

(r) **'Noce'**

cv. *Elepidote rhododendron*: 'Pineapple Delight' × unknown

**H:** J.-F. Petton (2001), **G:** J.-F. Petton (2015), **N:** J.-F. Petton (2018), **REG:** J.-F. Petton (2018)

Fls c.14/rounded truss, open funnel-shaped, 55 × 50mm, with 7 wavy-edged lobes, strong purplish pink (62A) in bud, opening inside light yellowish pink (19B); outside light yellowish pink (19B), with pale pink margins; stamens 12; style slightly curved, with a few hairs; ovary glabrous.

Calyx petaloid, the same length as the corolla, light yellowish pink (19B). Lvs elliptic, 110 × 45mm, asymmetrically rounded at base, obtuse at apex, margins decurved, hairless. Shrub 1.35 × 1.30m in 17 years. May. Hardy to -15C.

Nomenclatural standard: colour print provided by the registrant (WSY1015107). Etymology: named for the registrant's granddaughter.

(r) 'Pedro'

cv. Elepidote rhododendron: *yakushimanum*

'Koichiro Wada' (s) × *falconeri* (white form)

**H:** K. Heger (pre-1992), **G:** J. Jäck (2014), **N:** J. Jäck (2017), **REG:** J. Jäck (2018)

Fls up to 17/flattened ball truss, ventricose-campanulate, 50 × 60mm, with 7 emarginate, wavy-edged lobes, buds light pink, opening inside white, with a purplish red blotch at base; outside white. Lvs lanceolate, semi-glossy and green above, with ochre-coloured hairs below. Shrub 1.6 × 1.4m in 26 years. Mid April to early May. Hardy to -22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125516). Etymology: named for the brother of the exchange student of the registrant.

(r) 'Peg of My Heart'

cv. Elepidote rhododendron: *fortunei* subsp. *discolor*, open-pollinated (seed collected from the National Arboretum, Washington, D.C., USA, in the early 1980s)

**G:** T. Dove (2004), **S:** T. Dove (2018), **N:** T. Dove (2018), **REG:** T. Dove (2018)

Fls single, 10-14/domed truss, funnel-shaped, c. 50 × 80mm, with 7 rounded, shallowly toothed lobes, very light purple (76B) in bud, opening inside very pale purple (76C), shading to very pale purple (76D) at the centre of the lobes, with brilliant greenish yellow (151D) at throat and as spots along the dorsal lobe; outside very pale purple (76C) shading to very pale purple (76D); most stamens aborted; moderate, sweet and spicy scent. Calyx insignificant. Lvs elliptic, 190-220 × 33-57mm, rounded at base, broadly acute at apex, margins flat, semi-glossy, deep yellowish green (141A) above, strong yellowish green (141C) below, with sparse minute hairs on midrib (below) and on petioles. Shrub 4.0m × 3.7m in 33 years. Late season (early June in Washington, D.C., USA). Hardy to -25C (buds -22C).

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125536).

Etymology: named for the mother-in-law of the registrant.

(a) 'Pirin'

cv. Evergreen azalea: 'Mariko' (s) × 'Kermesinum'

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2-3/truss, funnel-shaped, 25 × 55mm, with 5-6 wavy-edged lobes, buds vivid purplish red (66A), with moderate purplish pink (68C) along ribs, opening inside vivid purplish red (57B), with vivid red

(57A) spots on the dorsal and adjacent lobes; outside vivid purplish red (57B-C); filaments strong purplish red (67A), with light purplish pink (55C) at base, anthers dark greyish purple (202A); style deep purplish pink (66C), stigma deep red (60A); not scented. Calyx 2mm long, strong yellow-green (144C), with long white hairs. Pedicels moderate reddish orange (178D), with touches of strong yellow-green (144C). Lvs elliptic, 35 × 15mm, cuneate at base, broadly acute at apex, with wavy margins, matt, moderate olive green (147A), with white hairs above, moderate yellow-green (147B), with brown hairs along midrib, below; petioles with very dark brown hairs. Shrub 0.5 × 0.8m in 10 years. Mid to late May (in northern Germany). Hardy to -24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125553).

Etymology: named after the Pirin National Park in Bulgaria.

(a) 'Purnululu'

cv. Evergreen azalea: 'Mariko' (s) × 'Kermesinum'

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2-3/truss, funnel-shaped, 21 × 53mm, with 5-6 wavy-edged lobes, buds vivid red (57A) and deep purplish pink (58D), opening inside vivid purplish red (57B), with vivid red (57A) spots on the dorsal and adjacent lobes; outside vivid purplish red (57B-C); filaments strong purplish red (67A), with light purplish pink (55C) at base, anthers dark greyish purple (202A); style deep purplish pink (66C), stigma deep red (60A); not scented. Calyx 2mm long, light yellow-green (145C), with strong greenish yellow (153C) at base. Pedicels moderate reddish orange (178D), with touches of strong yellow-green (144C). Leaves elliptic, 35 × 15mm, cuneate at base, broadly acute at apex, with wavy margins, matt, moderate olive green (147A), with white hairs above, moderate yellow-green (147B), with brown hairs along midrib, below; petioles with very dark brown hairs. Shrub 0.4 × 0.8m in 10 years. Mid to late May (in northern Germany). Hardy to -24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125554).

Etymology: named after the Purnululu National Park in Western Australia.

(r) 'RAF100'

cv. Elepidote rhododendron: un-named seedling (s) × 'Old Port'

**H:** C. Mugridge (2002), **G:** C. Mugridge (2018), **N:** C. Mugridge (2018), **I:** Millais Nurseries (2018), **REG:** C. Mugridge (2018)

Fls 16-20/ball truss, broadly funnel-shaped, 50 × 90mm, with 5 flat-edged lobes, buds deep mauve, opening inside very pale purple (75D), shading to light purple (75B) at margins, with moderate purplish red (58A) blotch and spots on dorsal lobe (the blotch sometimes split in two); outside light purple (75B); anthers cream; pistil yellow; not scented. Calyx 1-2mm long, green. Lvs elliptic, 160 × 50mm, cuneate at base, acute at apex, margins decurved, with a

satin-like surface, hairless. Shrub 1.0 × 1.0m in 16 years. Mid-season. Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125562). Etymology: named by the registrant, an ex-RAF pilot, to commemorate the centenary of the Royal Air Force in 2018.

(r) **'Reileyridge Lisa Marie'**

cv. *Elepidote rhododendron*: 'Scintillation' (s) × 'Janet Blair' × 'Crest'

**H:** H.E. Reiley (2007), **G:** M.L. Reiley (2016), **N:** M.L. Reiley (2016), **REG:** Reiley Ridge Nursery (2018)

Fls single, 7/truss, broadly funnel-shaped, 51 × 76mm, with 5 or 6, rounded, sometimes emarginate, wavy-edged lobes, strong purplish red (61B) in bud, opening inside pale purplish pink (62D) at margins, shading to pale greenish yellow (9D) at centre, with strong purplish red (63A) throat and strong purplish red (63B) spotting on the dorsal and adjacent lobes; outside pale purplish pink (62D) at margins and along midrib, shading to pale yellow (8D); stamens (12) and pistil yellowish. Calyx 6mm long, light yellow-green (145B). Lvs oblong, 102 × 51mm, rounded at base, obtuse at apex, margins flat, semi-glossy, moderate olive green (137A), hairless. Shrub 1.8 × 1.5m in 11 years. Mid-season (late May in Maryland, USA). Hardy to -18C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125502).

Etymology: named for a granddaughter of the hybridiser.

(r) **'Remember Rosemary'**

cv. *Elepidote rhododendron*: parentage unknown, but with characteristics of *williamsianum*

**H:** W.G.T. Hyde (1960s), **G:** W.G.T. Hyde (1970s), **N:** Mrs R.G. Legrand (2018), **I:** Mrs R.G. Legrand (2018), **REG:** Mrs R.G. Legrand (2018)

Fls 9–13/compact truss, open funnel-shaped, 53 × 30mm, with 5 very wavy-edged lobes, strong purplish red (54A) in bud, opening inside light purplish pink (62C), paler in the centre; outside moderate purplish pink (65A); with strong red (46A) spotting on the dorsal lobe. Calyx irregular, 2–4mm long, light yellow-green (145C). Lvs elliptic, 74 × 46mm, cordate at base, acute at apex, margins upcurved, matt green above, hairless. Shrub 2.6 × 1.8m in c.40 years. April.

Nomenclatural standard: colour print provided by the registrant (WSY0108685). Etymology: named for Rosemary Howell (an old friend of the registrant) and for all who share the name Rosemary.

(r) **'Rosea Fovea'**

cv. *Elepidote rhododendron*: 'Faggetter's Favourite' (s) × 'Kluis Sensation'

**H:** C. Mugridge (2001), **G:** C. Mugridge (2016), **N:** T. Mugridge (2016), **I:** Millais Nurseries (2018), **REG:** C. Mugridge (2018)

Fls 17–20/ball truss, broadly funnel-shaped, 50 × 90mm, with 6 wavy-edged lobes, buds deep red, opening inside pale purplish pink (56B), with deep

pink (52C) at rim, lightly spotted with strong red (53B) at the base of the dorsal lobe; outside pale purplish pink (56B), with deep pink (52C) at rim; anthers brown; pistil yellow; not scented. Calyx 1–2mm long, pale green. Lvs elliptic, 150 × 55mm, rounded at base, broadly acute at apex, margins flat, matt above, hairless. Shrub 1.0 × 1.0m in 17 years. Mid-season. Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125563).

Etymology: named, by the son of the registrant (an optometrist), after part of the eye.

(r) **'Rose Passion'**

cv. *Elepidote rhododendron*: 'Summer Sunrise' (s) × 'Plum Passion'

**H:** J. Barlup (2009), **G:** J. Barlup (2014), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 17/ball truss, broadly funnel-shaped, 44 × 70mm, with 5 rounded, wavy-edged lobes, strong red (53B) in bud, opening inside pale yellow (11D), shading through pale purplish pink (55D) to strong purplish pink (55B) at margins, and with darker midveins and an area of deep red (53A) at the base, extending 13mm dorsally, changing to strong red (53D) spotting on the dorsal lobe; outside pale purplish pink (55D), with strong purplish pink (55B) midveins towards the margins; filaments reddish; style reddish, with a darker red stigma; not scented. Calyx 3mm long, pale yellow (11D), with a deep red (53A) flare. Lvs elliptic, 114 × 44mm, rounded at base, broadly acute at apex, margins flat, moderate olive green (147A) above, hairless. Shrub 0.4 × 0.6m in 7 years. Mid-season (early May in Seattle area, Washington, USA). Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108651).

(r) **'Rose Velvet'**

cv. *Elepidote rhododendron*: 'Cody' (s) × 'Plum Passion'

**H:** J. Barlup (2009), **G:** J. Barlup (2014), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 17/domed truss, broadly funnel-shaped, 57 × 83mm, with 5–6, rounded, wavy-edged lobes, strong red (53C) in bud, opening inside strong purplish red (55B), shading near base to pale yellow (11D), with a strong red (53D) dorsal flare from the base, becoming spots, and with small areas of greenish white (155C) near the base producing a central star-like effect; outside strong purplish red (55B), with deep purplish pink (55A) margins and midveins; filaments pinkish; style pinkish, with a green stigma. Calyx 13mm long, pale yellow (11D), with strong red (53D) streaking. Lvs elliptic, 127 × 51mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.8m in 8 years. Mid-season (mid May in Seattle area, Washington, USA). Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108653).

(r) **'Sarah's Song'**

cv. *Elepidote rhododendron*: un-named seedling (s) × 'Lem's Cameo'

**H:** C. Mugridge (2001), **G:** C. Mugridge (2017),  
**N:** C. Mugridge (2018), **I:** Millais Nurseries (2018),  
**REG:** C. Mugridge (2018)

Fls 15–17/ball truss, broadly funnel-shaped, 35 × 95mm, with 6–7 wavy-edged, ruffled lobes, buds deep pink, opening inside light purplish pink (62C), shading to deep purplish pink (64D) at margins, with deep purplish red (59B) blotch and spots on the dorsal lobe; outside light purplish pink (62C), shading to deep purplish pink (64D) at margins; anthers cream; pistil green. Normal calyx absent. Lvs elliptic, 150 × 55mm, rounded at base, broadly acute at apex, margins flat, with a satin-like surface, hairless. Shrub 1.0 × 1.0m in 17 years. Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125564).

Etmymology: named for the daughter of the registrant.

(r) **'Savanna Gold'**

cv. *Elepidote rhododendron*: 'Hachgold' (GOLDEN EVEREST) (s) × 'Olga'

**H:** Holger Hachmann (2005), **G:** Baumschule H. Hachmann, **N:** Holger Hachmann, **I:** Baumschule H. Hachmann, **REG:** O. Tschetsch (2018)

Fls 11–14/domed truss, funnel-shaped, 30 × 70mm, with 5 wavy-edged lobes, buds vivid reddish orange (34A/42A), opening inside vivid yellow (13A), with deep yellowish pink (39B) margins, becoming brilliant yellow (13C), with vivid yellow (13A) along the midrib and with spots of deep red (53A) and strong red (53B) on the dorsal and adjacent lobes; outside brilliant yellow (13B/C), with strong red (39A) at margins, fading with maturity; 10 stamens, filaments pale greenish yellow (2D), anthers brownish orange (165B); style brilliant yellow-green (150C), stigma strong orange (26A); ovary strong yellowish green (141C), with short white hairs; not scented. Calyx 4mm long, entire, brilliant yellow (13C), with short dark brown hairs. Pedicels 30mm long, brilliant yellow-green (149B), with moderate red (180B). Lvs elliptic, 95 × 40mm, rounded at base and apex, with upcurved margins, matt, moderate olive green (147A) above, strong yellow-green (143C) below, hairless; petioles 25mm long, strong yellow-green (144B), with moderate red (181A) above and strong yellow-green (144B), with deep yellowish pink (39B) below. Shrub 1.1 × 1.4m in 10 years. Mid to late May (in northern Germany). Hardy to –22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125611).

(a) **'Schnupper's Uluru'**

cv. *Evergreen azalea*: 'Mariko' (s) × 'Kermesinum'

**H:** F. Schnupper (2003), **G:** F. Schnupper,  
**N:** F. Schnupper (2018), **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2–3/truss, funnel-shaped, 20 × 50mm, with 5–6 wavy-edged lobes, buds vivid purplish red (66A), opening inside vivid purplish red (66B) with deep red (60A) spots on the dorsal and adjacent

lobes; outside strong purplish red (64B); filaments strong purplish red/strong purplish pink (67A/D), anthers dark greyish purple (202A); style vivid purplish red/deep purplish pink (67B/C), with moderate reddish orange (179C) at base, stigma moderate reddish orange (179C); not scented. Calyx 2mm, strong yellow-green (144C), with long white hairs. Pedicels light yellow-green (144D), with moderate red (180A). Lvs obovate, 30 × 14mm, cuneate at base, broadly acute at apex, margins wavy, matt, moderate olive green (147A), with white hairs above, moderate yellow-green (146B), midrib brownish orange (165B), with long brown hairs below; petioles 5mm long. Shrub 0.4 × 0.8m in 10 years. Mid May to early June (in northern Germany). Hardy to –24C.

Trade designation: ULURU

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125555).

Etmymology: named after the Uluru-Kata Tjuta National Park in the Northern Territory, Australia.

(a) **'Scotian Fire'**

cv. *Evergreen azalea*: 'Pink Clusters' (s) × 'Mount Seven Star'

**H:** J.K. Weagle (1983), **G:** J.K. Weagle (1985),  
**N:** J.K. Weagle (1990), **I:** Rhodo Land Nursery (c.2003), **REG:** J.K. Weagle (2018)

Fls single, 2/terminal truss, open funnel-shaped, 35 × 47mm, with 5 flat lobes, broadly acute at apex, buds vivid red to vivid reddish orange (44A–44B), opening inside vivid red to vivid reddish orange (44A–44B), with barely discernible blotch and spots of vivid red (45A) on the dorsal lobe; outside vivid red to vivid reddish orange (44A–44B), very glossy at base; stamens and pistil red; not scented. Calyx 8mm long, brilliant yellow-green (149A). Leaves elliptic, 35 × 17mm (summer), 17 × 10mm (winter), cuneate at base, broadly acute at apex, with flat margins, glossy moderate olive green (146A) above, moderately covered with adpressed silver white hair below, with midrib; petioles and new shoots heavily covered. Shrub 0.6 × 0.6m in 10 years. Late (mid to late June in coastal Nova Scotia, Canada). Hardy to –25C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125537).

(r) **'Shannon for You'**

cv. *Elepidote rhododendron*: parentage unknown

**H:** H.E. Reiley (2006), **G:** M.L. Reiley (2016),  
**S:** M.L. Reiley (2017), **N:** M.L. Reiley (2018),  
**REG:** Reiley Ridge Nursery (2018)

Fls single, 13/truss, broadly funnel-shaped, 57 × 90mm, with 6 rounded, very wavy-edged, reflexed and partially emarginated lobes, strong reddish purple (78B) in bud, opening inside creamy white, with strong purple (77B) picotee and with vivid yellow-green (154A) at throat; outside strong purple (77B), blended with light purple (77C); filaments white; style greenish yellow, stigma greenish yellow, touched purple; lightly scented. Lvs elliptic, 140 × 51mm, rounded at base, broadly acute at apex, with margins downcurved, semi-glossy, moderate olive green (137B), hairless. Shrub 0.9 × 1.0m in 12 years. Mid-season (May in

Maryland, USA). Hardy to -15C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125503).  
Etymology: named for the granddaughter of the hybridiser.

(r) ‘**Sievershütten**’

cv. Elepidote rhododendron: ‘Scarlet Wonder’ (s) × ‘Goldkrone’

**H:** J. Jäck (1992), **G:** J. Jäck (1998), **N:** J. Jäck (1998), **REG:** J. Jäck (2018)

Fls single, 7–8/loose, open-topped truss, funnel-shaped, 45–55 × 55–65mm, with 5 wavy-edged lobes, buds dark red-pink, opening inside dark red-pink, with a slight blotch of dark red spots on the dorsal lobe; outside dark red-pink. Lvs elliptic, 70 × 40–45mm, obtuse at base, rounded at apex, with flat margins, matt green, hairless. Shrub 0.75 × 0.80m in 25 years. Early May. Hardy to -20C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125519).

Etymology: named after a town near Hamburg, Germany.

(r) ‘**Sir Paul Jenkins**’

cv. Elepidote rhododendron: *praestans* (s) × unknown

**H:** Exbury Gardens, **G:** Exbury Gardens, **N:** Lionel de Rothschild (2018), **REG:** T. Clarke (2018)

Fls 23/truss (the truss compact and rounded at first, but becoming more lax), ventricose-campanulate, 55 × 55mm, with 8 slightly wavy-edged lobes, inside very pale purple (75D) fading to very pale purple (69D), and with a stripe below the lobe sinuses of light purple (75A), which fades to very pale purple (69B), with a dark red (59A) blotch at the base of the dorsal throat; outside similarly coloured; filaments white, anthers brown; stigma prominent, lime green. Calyx insignificant, light yellow-green (145B). Lvs oblanceolate, 240 × 95mm, obliquely cuneate at base, obtuse at apex, margins flat, matt green above, with an agglutinated indumentum below of moderate orange-yellow (164B) when young, which matures to pale yellow (160D). Shrub 5.0 × 5.0m in 15 years. April. Hardy to -7C.

Nomenclatural standard: colour print provided by the registrant (WSY0108686). Etymology: named for a friend of the nominant.

(r) ‘**Sonora Sun**’

cv. Elepidote rhododendron: ‘Maverick’ (s) × ‘Trina’

**H:** J. Barlup (2009), **G:** J. Barlup (2014), **N:** J. Barlup (2017), **REG:** J. Barlup (2018)

Fls 17/ball truss, broadly funnel-shaped, 51 × 83mm, with 5 rounded, emarginate and wavy-edged lobes, moderate red (47A) in bud, opening inside light yellow (18B), shading to pale yellow (18D), with strong red (53C) dorsal spotting; outside light yellow (18B), shading to pale yellow (18D), with moderate yellowish pink (31D) midveins; filaments yellow; style yellow, with a brown stigma; not scented. Calyx insignificant. Lvs elliptic, 108 × 41mm, rounded at base, broadly acute at apex, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.8m in 8 years. Mid-season (early May in

the Seattle area, Washington, USA). Hardy to -15C.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108656).

(r) ‘**Stadt Lucianópolis**’

cv. Elepidote rhododendron: *yakushimanum*

‘Koichiro Wada’ (s) × *falconeri* (white form)

**H:** K. Heger (1992), **G:** J. Jäck (2010), **N:** J. Jäck, **REG:** J. Jäck (2018)

Fls single, 20/dense, rounded truss, campanulate, 55 × 65mm, with 6–7 wavy-edged lobes, buds pale pink, with touches of darker red-pink, opening inside white, with dark red-pink blotch at base; outside white. Lvs lanceolate 140–165 × 50–65mm, obtuse to rounded at the base, acute at apex, with margins a little decurved, matt green above, with silvery white hairs when young, aging to a thick, beige indumentum. Shrub 1.6 × 1.1m in 28 years. Late April to early May. Hardy to -22C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125522).

Etymology: named after the town in São Paulo, Brazil.

(a) ‘**Talamanca**’

cv. Evergreen azalea: ‘Mariko’ (s) × ‘Madame Albert van Hecke’

**H:** F. Schnupper (2003), **G:** F. Schnupper, **N:** F. Schnupper, **I:** Schnupper Baumschule (2018), **REG:** O. Tschetsch (2018)

Fls single, 2–3/truss, funnel-shaped, 25 × 45mm, with 5–6 lobes, with flat margins, buds vivid purplish red (61C) and moderate purplish pink (62B), opening inside vivid purplish red (67B), with moderate purplish red (59C) spots on the dorsal and adjacent lobes; outside vivid purplish red (67B); filaments vivid purplish red (61C) and strong purplish pink (62A), with short white hairs on the basal half, anthers dark red (187A); style deep pink (185D), with light yellow-green (144D) at base, stigma strong greenish yellow (151A); not scented. Calyx 2mm long, strong yellow-green (141D). Pedicels 10mm long, moderate reddish orange (171A) and strong yellow-green (141D), with white hairs. Lvs elliptic, 31 × 13mm, cuneate at base, broadly acute at apex, with flat margins, glossy, moderate olive green (147A), with sparse white hairs above, moderate olive green (146A), fading to moderate yellow-green (146B) below. Shrub 0.3 × 0.8m, in 10 years. Mid May to early June (in northern Germany). Hardy to -24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125556).

Etymology: named after the Cordillera de Talamanca mountain range in La Amistad National Park on the Costa Rica-Panama border.

(r) ‘**Tante Jenny**’

cv. Elepidote rhododendron: *rex* hybrid (from seed collected from Hamburg Botanic Garden in 1992)

**G:** J. Jäck (2010), **N:** J. Jäck (2017), **REG:** J. Jäck (2018)

Fls single, 18–20/roundish, flattened, somewhat loose truss, funnel-shaped, 65 × 80mm, with 7 wavy-edged lobes, buds bright, deep lilac pink, opening inside white, with a solid dorsal blotch of deep red and with deep red spots radiating along the

dorsal lobe; outside white. Lvs lanceolate, 200 × 70mm, obtuse at base, acute at apex, with flat margins, semi-glossy, green above, with silvery white, becoming brownish, hairs below. Shrub 2.5 × 2.0m in 25 years. Mid April to early May. Hardy to -22C. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125526). Etymology: named for a favourite aunt of the registrant.

(a) **'Trouscoat Daoulagad'**

cv. Deciduous azalea: 'Whitethroat' (s) × 'Sunte Nectarine'

**H:** L. Le Duigou (2004), **G:** A. Le Duigou (2010), **N:** A. & L. Le Duigou (2018), **I:** Les Pépinières Nicolandes (2018), **REG:** A. & L. Le Duigou (2018) Fls 9/ball truss, tubular funnel-shaped, 50 × 60mm, with 5 wavy-edged lobes, deep pink (52B) in bud, opening inside in some flowers strong pink (54C), fading to pale purplish pink (56B), in others white (NN155D), with a brilliant yellow (20A) blotch; outside with a strong purplish red (54A) tube and strong purplish pink (55B) lobes; brilliant yellow (20A) spots on the dorsal lobe; stamens 5-7; style pale purplish pink (55D), with a strong yellow-green (144A) stigma; strongly scented. Calyx 5mm long, brilliant yellow-green (150A). Lvs elliptic, 63 × 26mm, cuneate at base, broadly acute at apex, margins flat, hairless. Shrub 1.3 × 0.9m in 14 years. May. Hardy to -15C. Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0125650). Etymology: the registrants' botanical garden is called Les Jardins de Trouscoat, whilst "Daoulagad" means "two eyes" in Breton.

(r) **'Trouscoat Flourdiliz'**

cv. Elepidote rhododendron: 'Lodauric Iceberg' (s) × Fred Wynniatt Group

**H:** L. Le Duigou (2003), **G:** A. Le Duigou (2012), **N:** A. & L. Le Duigou (2018), **I:** Les Pépinières Nicolandes (2018), **REG:** A. & L. Le Duigou (2018) Fls 11/lax truss, narrowly funnel-shaped, 75 × 95mm, with 7 wavy-edged lobes, deep purplish pink (58D) in bud, opening inside pale purplish pink (62D), paling towards the centre; outside light purplish pink (62C), shading to pale yellow (11D) towards the base; unmarked; stamens 13; style 60mm long, more-or-less straight, light greenish yellow (1C), with a strong yellow-green (143C) stigma; scented. Calyx 5mm long, strong yellow-green (N144C), with a margin of vivid purplish red (N57A). Lvs oblong, 180 × 80mm, rounded at base, obtuse at apex, margins flat, hairless. Shrub 2.25 × 2.10m in 15 years. May to June. Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0125655). Etymology: the registrants' botanical garden is called Les Jardins de Trouscoat, whilst "Flourdiliz" means "lily" in Breton.

(r) **'Trouscoat Gwagennek'**

cv. Elepidote rhododendron: 'Double Date' (s) × 'Anatevka'

**H:** L. Le Duigou (2010), **G:** A. Le Duigou (2015), **N:** A. & L. Le Duigou (2018), **I:** Les Pépinières Nicolandes (2018), **REG:** A. & L. Le Duigou (2018)

Fls 13/conical truss, saucer-shaped, 60 × 95mm, with 7 strongly wavy-edged lobes, strong purplish red (72A) in bud, opening inside light purple (75A), fading to centre, with short, petaloid stamens which are strong reddish purple (72B); outside light purple (75A), with strong reddish purple (72B) ribs; dorsal lobe with dark red (59A) spots, other lobes with strong purplish red to strong purplish pink (63B-C) spots, the spots showing through to the outside; style curved, pale greenish yellow (1D), stigma moderate red (182A). Calyx 4mm long, strong yellow-green (144B). Lvs elliptic, 140 × 40mm, cuneate at base, acute at apex, margins flat, hairless. Shrub 1.1 × 1.1m in 7 years. May.

Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0125659).

Etymology: the registrants' botanical garden is called Les Jardins de Trouscoat, whilst "Gwagennek" means "wavy" in Breton.

(r) **'Trouscoat Roz'**

cv. Elepidote rhododendron: 'Bluebell' (s) × 'One Thousand Butterflies'

**H:** L. Le Duigou (2006), **G:** A. Le Duigou (2012), **N:** A. & L. Le Duigou (2018), **I:** Les Pépinières Nicolandes (2018), **REG:** A. & L. Le Duigou (2018) Fls 19/domed truss, open funnel-shaped, 50 × 95mm, with 5 wavy-edged lobes, strong purplish red (71B) in bud, opening inside very light purple (75C), with deep purplish pink (N74C) margins and with an area of deep red (53A) at the base of the throat; outside vivid purplish red (67B); dorsal lobes with moderate red (179B) markings; stamens 9; style curved, deep purplish pink (67C), with a strong yellow-green (144B) stigma. Calyx 2-5mm long, strong yellow-green (144A). Lvs elliptic, 160 × 45mm, cuneate at base, broadly acute at apex, margins flat, hairless. Shrub 1.1 × 1.4m in 11 years. May. Hardy to -15C.

Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0125663).

Etymology: the registrants' botanical garden is called Les Jardins de Trouscoat, whilst "Roz" means "Pink" in Breton.

(r) **'Tricia Cox'**

cv. Lepidote rhododendron: selection from *formosum* (from Cox & Hutchison 301) (s) × selection from *dendrocharis* (from the wild collection made as CCH 3915)

**I:** Glendoick Gardens Ltd, **REG:** Glendoick Gardens Ltd (2018)

Fls 2-3/truss, openly funnel-shaped, 50 × 60mm, with 5 wavy-edged, slightly ruffled lobes, buds deep purplish pink (54B), opening inside pale purplish pink (56A-56D) and pale yellowish pink (36D), with pinkish red spots in the throat; outside pale purplish pink (56C), flushed with deeper purplish pink shades (54B/55A); scented. Calyx 3-6mm long, green. Lvs ovate, 15-30 × 7-12mm, ovate at base, mucronate at apex, margins flat to decurved, glossy above, with scales. Shrub 0.2 × 0.3m. April to May. Hardy to -10C.

Etymology: named for the wife of one of the hybridisers.

(r) 'Twenty Four Carat'

Syn. of '24 Karat'

(a) ULURU

Trade designation for 'Schnupper's Uluru'

(r) 'Vanillefeuer'

cv. Elepidote rhododendron: *smirnowii* (s) × *dichroanthum* subsp. *scybocalyx* × *wardii*

H: H.L. Meltzer (1991), G: J. Jäck (1996), N: J. Jäck, REG: J. Jäck (2018)

Fls single, 8–10/dense, slightly flattened truss, funnel-shaped, 50 × 70–80mm, with 5 wavy-edged lobes, buds deep pink, opening inside creamy white, with a small deep reddish pink blotch at the base of the dorsal lobe and with a flare of deep pinkish red spots radiating out towards the margins; outside creamy white. Lvs oblanceolate, 90–140 × 35–45mm, obtuse at base, acute and mucronate at apex, margins flat, matt green above, with white hairs, becoming silvery below. Shrub 2.5 × 2.5m in 27 years. Early to mid May. Hardy to –26C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125529).

Etymology: named for the colour of the flowers and the flame-like markings within them.

(a) 'Vidzeme'

cv. Deciduous azalea: 'Homebush' (s) × 'Austra'

H: R. Kondratovičs (1999), G: R. Kondratovičs (2006), N: G. Riekstiņa (2018), REG: Rhododendron Nursery Babite, University of Latvia (2018)

Fls double, 15/strongly domed truss, tubular funnel-shaped, 40 × 50mm, with 2 whorls of 5 flat lobes, buds moderate red (N34C), opening inside strong purplish pink (62A), with pinkish white (N155B) at midrib, dorsal and adjoining lobes marked with a vivid yellow (15A) blotch; outside pink, with pinkish white (N155B) at midrib; style orange, stigma green; lightly scented. Calyx 1mm long, green. Lvs elliptic, 80 × 35mm, acute at base and apex, with wavy margins, glossy above, hairless. Shrub 2.2 × 2.4m in 18 years. Late May to early June. Hardy to –27C.

Nomenclatural standard: colour print provided by the registrant (WSY0125575). Etymology: named after one of the Latvian historical and cultural regions.

(r) 'Walter Eckel'

cv. Elepidote rhododendron: 'Bambi' (s) × *falconeri* (white form)

H: K. Heger (1992), G: J. Jäck (2000), N: J. Jäck (2016), REG: J. Jäck (2018)

Fls single, c.11/loose, open-topped truss, campanulate, 40 × 60mm, with 6 wavy-edged lobes, buds deep pinkish red, opening inside paler pinkish red, with touches of a deeper tone at the lobe margins and with a deeper purplish red blotch at base; outside paler pinkish red. Calyx c. 5mm long, pale pinkish red, with touches of a yellow tone. Lvs elliptic, 70–120 × 30–50mm, obtuse at base, broadly acute at apex, glossy green above, with beige hairs below. Shrub 1.2 × 1.0m in 24 years. Early May. Hardy to –18C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125533).

Etymology: named for a former teacher of the registrant.

(r) 'Whimbrel'

cv. Lepidote rhododendron: *calostrotum* 'Gigha' (s) × 'Plover'

I: Glendoick Gardens Ltd, REG: Glendoick Gardens Ltd (2018)

Fls 1–3/truss, funnel-shaped, very flat-faced, 30 × 45mm, with 5 wavy-edged lobes, buds vivid purplish red (67B), opening inside strong purplish pink (67D), overlaid with a deeper tone (67C) towards apex, with vivid purplish red (67B) spots on the dorsal and adjoining lobes; outside strong purplish pink (67D), overlaid with a deeper tone (67C) towards apex; filaments pale, with black anthers; style strong purplish red (67A), with black stigma scented. Calyx 5mm long, green, touched purplish pink at margins. Lvs elliptic, 32 × 20mm, obtuse at base and apex, margins flat to decurved, glossy above, with scales. Mid-season (April to May in Scotland). Hardy to –18C.

(r) 'Whispering Giant'

cv. Elepidote rhododendron: 'Lem's Cameo' (s) × 'Faggetter's Favourite'

H: C. Mugridge (2002), G: C. Mugridge (2018), N: C. Mugridge (2018), I: Millais Nurseries (2018), REG: C. Mugridge (2018)

Fls 16–21/conical truss, broadly funnel-shaped, 40 × 100mm, with 5 wavy-edged, slightly ruffled lobes, buds deep pink, opening inside very pale purple (73D), washed with strong purplish pink (63C) at margins and with a slightly deeper tone around the very edge of the lobes, dorsal lobe spotted with strong purplish red (63A) from base to two-thirds of the way up the lobe; outside very pale purple (73D), washed with strong purplish pink (63C) at margins; anthers brown; pistil red; not scented. Calyx 1–3mm long, pale green. Lvs elliptic, 150 × 60mm, rounded at base, broadly acute at apex, margins flat, matt above, hairless. Shrub 1.0 × 1.0m in 12 years. Mid-season. Hardy to –15C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125565).

Etymology: named after the Bristol Britannia aircraft, that was the last aircraft captained by the registrant for the RAF.

(r) 'White June'

cv. Elepidote rhododendron: *yakushimanum* (s) × *maximum*

G: Baumschule H. Hachmann, N: Holger Hachmann, I: Baumschule H. Hachmann, REG: Ö. Tschetsch (2018)

Fls 15–18/ball truss, funnel-shaped, 35 × 55mm, with 5 wavy-edged lobes, buds pale purplish pink (62D), marked with strong purplish pink (62A), opening inside yellowish white (155D), with strong greenish yellow (151B) spots on the dorsal lobe; outside yellowish white (155D); 10 stamens, filaments yellowish white (155D), with dense white hairs at base, anthers moderate yellow (162A–B); style greenish white (157D), stigma greyed red (179C–D); ovary with dense white hairs; not scented. Calyx 2mm long, light yellow-green (150D), translucent. Pedicels 40mm long, brilliant yellow-green (150C).

Lvs elliptic, 120 × 40mm, rounded at base, mucronate at apex, with decurved margins, matt, moderate olive green (147A) with dense white hairs on opening, thinning, but never going, especially on the margins, where they become grey white above, with dense moderate yellow (161A) hairs below; petioles strong yellow-green (144C). Shrub 0.9 × 1.2m in 10 years. Late May to mid June (in northern Germany). Hardy to -24C.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125614). Possibly from seed obtained from the American Rhododendron Society seed exchange.

(r) **'Yellow Bliss'**

cv. Elepidote rhododendron: 'Hotei' (s) × 'Wickiup'

**H:** H.R. Helm (1990), **G:** D. Denkers,

**N:** H.R. Helm (2018), **REG:** H.R. Helm (2018)

Fls single, 15/ball-shaped truss, funnel-shaped, 44 × 51mm, with 5 rounded, often emarginate lobes, with flat margins, vivid red (45C) in bud, opening inside orange, quickly becoming vivid yellow (12A), with vivid red (45C) spots on the dorsal and adjoining lobes, spots fading to vivid yellow (12A); outside vivid yellow (12A), with vivid red (45C) spots, spots fading to vivid yellow (12A); filaments yellow, anthers brown; style yellow, with a yellow-brown stigma; not scented. Calyx 25mm long, vivid yellow (12A), with vivid red (45C) overlay, particularly on dorsal side, the red fading to yellow on maturity. Lvs oblanceolate, 76–102 × 44mm, cuneate at base, broadly acute at apex, margins flat, semi-glossy and brilliant yellowish green (140B) above, hairless. Shrub 1.8 × 2.4m in 28 years. Mid-season (late May in Seattle, Washington, USA). Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0125499).

(a) **'Zemgale'**

cv. Deciduous azalea: 'Homebush' (s) × 'Austra'

**H:** R. Kondratovičs (1999), **G:** R. Kondratovičs (2006), **N:** G. Riekstiņa (2018),

**REG:** Rhododendron Nursery Babīte, University of Latvia (2018)

Fls double, 16/slightly domed truss, tubular funnel-shaped, 35 × 40mm, with 2 whorls of 5 wavy-edged lobes, buds moderate reddish orange (35B), opening inside strong yellowish pink (32C); outside light orange-yellow (24C); style orange-red, stigma green; lightly scented. Calyx 2mm long, green, Lvs elliptic, 70 × 35mm, acute at base and apex, with wavy margins, glossy green above, hairless. Shrub 2.4 × 2.4m in 18 years. Late May to early June. Hardy to -27C.

Nomenclatural standard: colour print provided by the registrant (WSY0125576). Etymology: named after one of the Latvian historical and cultural regions.

## Corrections to the *International Rhododendron Register and Checklist (2004)*

The following list indicates further significant additions and corrections that need to be made to the printed text of the Register and Checklist published in 2004 or to its subsequent Supplements. The Registrar is grateful to all those who have brought such errors to her notice and welcomes knowledge of further amendments that may need to be made. Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the Register and Checklist (or one of its Supplements) on which the entry concerned is to be found.

**'Barbar'** (p.108)

The correct spelling is 'Babar'

**'Birthday Girl'** (p.144)

Add AM 2000

**'Brimble'** (Fourth Supplement, p.3)

Add AM 2009

**'Charles Michael'** (p.231)

Add PC 2010

**'Crosswater Belle'** (p.300)

Add AM 2002

**'David Rockefeller'** (p.318)

Add PC 2005

**'Elfin Gold'** (p.399)

Add AM 2010

**'Fantastica'** (p.437)

Add AM 2004

**'Florida Ogada'** (p.459)

Add FCC 2002

**'Gle002'** (p.515 & Fourth Supplement, p.23)

Add AM 2009

**'Hampshire Belle'** (p.570)

Add AM 2009

**'Kiwi Majic'** (p.728)

The correct spelling is 'Kiwi Magic'

**'Lem's Monarch'** (p.776)

Add FCC 2002

**'Loch Awe'** (First Supplement, p.23)

Add AM 2010

**'Penvergate'** (Third Supplement, p.18)

Add AM 2009

**'Princess Margaret of Windsor'** (p.1107)

Add AM 2005

**'Tinner's Blush'** (Third Supplement, p.24)

Add AM 2010

**'Tower Dexter'** (p.1380)

Add AM 2000

**'Westport Point'** (p.1443)

Add AM 2009

**'Wisley Pearl'** (First Supplement, p.39)

Add AM 2004

## Names and Addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

**Barlup, J.**

Bellevue, Washington, USA

**Baumschule H. Hachmann**

Barmstedt, Germany

**Begg, L.**

Gembrooke, Victoria, Australia

**Blackmore, R.**

Victoria, British Columbia, Canada

**Borsky, M.**

Bridgewater, New Jersey, USA

**Brindabella Country Gardens**

Toowoomba, Queensland, Australia

**Bush, V.**

Huntsville, Alabama, USA

**Clagett, W.L.**

Derwood, Maryland, USA

**Clarke, T.**

c/o Exbury Gardens, q.v.

**Coble, S.**

Gettysburg, Pennsylvania, USA

**Denkers, D.**

Bainbridge Island, Washington, USA

**Dove, T.**

Harwood, Maryland, USA

**Ebbinghaus, R.**

Ahrensburg, Germany

**Exbury Gardens**

Hampshire, England

**Fosse, O.**

Bryne, Norway

**Furman, R.A.**

Brewster, Massachusetts, USA

**Glendoick Gardens Ltd**

Perth, Scotland

**Gray, J.C.**

c/o Brindabella Country Gardens, q.v.

**Greer, H.E.**

Eugene, Oregon, USA

**Haag, C.R. & V.**

Brevard, North Carolina, USA

**Hachmann, Holger**

c/o Baumschule H. Hachmann, q.v.

**Heger, Dr K.**

Krefeld, Germany

**Helm, H.R.**

Bainbridge Island, Washington, USA

**Hesse, H.**

Hamersen, Germany

**Humphries, K.F.**

White House Beach, Massachusetts, USA

**Huntsville Madison County Botanical Garden Society, Inc.**

Huntsville, Alabama, USA

**Hyde, W.G.T.**

Ferndown, Dorset, England

**Inkarho GmbH**

Bad Zwischenahn, Germany

**Jäck, J.**

Hamburg, Germany

**Jørgensen, P.M.**

Det Norske Arboret, Mildevegen, Hjellesstad, Norway

**Knights, T.**

USA

**Kondratovičs, R.K.**

Rīga, Latvia

**Le Duigou, A. & L.**

Scaër, France

**Legrand, R.G.**

West Moors, Dorset, England

**Les Pépinières Nicolandes**

Riec-sur-Bélon, France

**Manion, D.**

Eugene, Oregon, USA

**Meltzer, H.L.**

Neustadt, Bavaria, Germany

**Millais, D. & E.G.**

Millais Nurseries, q.v.

**Millais Nurseries**

Farnham, Surrey, England

**Mills, M.M.**

Philadelphia, Pennsylvania, USA

**Mugridge, C., Christopher & T.**

St Asaph, Denbighshire, Wales

**Murcott, R.**

East Norwich, New York, USA

**Petton, J.-F.**

Scaër, France

**Phipps Jr Estate, Howard**

Garrison, New York, USA

**Player, L.M.**

Fayetteville, North Carolina, USA

**Preil, W.**  
Ahrensburg, Germany

**Rabideau, R.R.**  
Camden, New Jersey, USA

**RareFind Nursery**  
Jackson, New Jersey, USA

**Reiley, H.E., M.L. & S.L.**  
c/o Reiley Ridge Nursery, q.v.

**Reiley Ridge Nursery**  
Woodsboro, Maryland, USA

**Rhododendron Nursery Babite, University of Latvia**  
Rīga, Latvia

**Rhodo Land Nursery**  
Niagra-on-the-Lake, Ontario, Canada

**Riekstiņa, G.**  
c/o Rhododendron Nursery Babite, q.v.

**Rothschild, E., Lionel & N. de**  
c/o Exbury Gardens, q.v.

**Schnupper, F.**  
c/o Schnupper Baumschule, q.v.

**Schnupper Baumschule**  
Westerstede, Germany

**Somerville, E.**  
Marietta, Georgia, USA

**Tschetsch, O.**  
Gabstede, Germany

**Vorwerk Pflanzenhandel GmbH**  
Rastede, Germany

**Weagle, J.K.**  
Halifax, Nova Scotia, Canada

**White's Nursery**  
Germantown, Maryland, USA

**Wickhorst, B.**  
c/o Baumschulen H. Hachmann, q.v.

**Woodard, G.**  
Westbury, New York, USA



ISBN 9781907057953



9 781907 057953